

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 0 984 021 A2

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:
08.03.2000 Bulletin 2000/10

(51) Int Cl.7: C07H 21/00

(21) Application number: 99306168.8

(22) Date of filing: 03.08.1999

(84) Designated Contracting States:
AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE
Designated Extension States:
AL LT LV MK RO SI

(30) Priority: 03.08.1998 US 128052
22.06.1999 US 338179

(71) Applicant: Hewlett-Packard Company
Palo Alto, California 94304 (US)

(72) Inventors:
• Dellinger, Douglas J.
Sunnyvale, CA 94087 (US)
• Caruthers, Marvin H.
Boulder, CO 80303 (US)
• Betley, Jason R.
Bury St. Edmunds, Suffolk IP28 6HW (GB)

(74) Representative: Tollett, Ian et al
Williams, Powell & Associates,
4 St. Paul's Churchyard Inn
London EC4M 8AY (GB)


(54) Synthesis of oligonucleotides

(57) The invention provides a method for synthesizing oligonucleotides using carbonate protection of hydroxyl groups and nucleophilic deprotection reagents. The deprotection reagents irreversibly cleave the carbonate protecting groups while simultaneously oxidizing the internucleotide phosphite triester linkage, and can be used in aqueous solution at neutral to mildly basic pH. The method eliminates the need for separate de-

protection and oxidation steps, and, since the use of acid to remove protecting groups is unnecessary, acid-induced depurination is avoided. Fluorescent or other readily detectable carbonate protecting groups can be used, enabling monitoring of individual reaction steps during oligonucleotide synthesis. The invention is particularly useful in the highly parallel, microscale synthesis of oligonucleotides. Reagents and kits for carrying out the aforementioned method are provided as well.

Figure 1

3'-5' Synthesis With DMT Protection



EP 0 984 021 A2

Description

[0001] This is a continuation-in-part of U.S. Patent Application Serial No. 09/128,052, filed August 3, 1998, the disclosure of which is incorporated by reference herein.

[0002] The present invention relates generally to nucleic acid chemistry and to the chemical synthesis of oligonucleotides. More particularly, the invention relates to an improved method for synthesizing oligonucleotides wherein carbonates are used as hydroxyl-protecting groups and "alpha-effect" nucleophiles such as peroxides are used in the deprotection thereof. The invention has utility in the fields of biochemistry, molecular biology and pharmacology, and in medical diagnostic and screening technologies.

[0003] Solid phase chemical synthesis of DNA fragments is routinely performed using protected nucleoside phosphoramidites. S.L. Beaucage *et al.* (1981) *Tetrahedron Lett.* **22**:1859. In this approach, the 3'-hydroxyl group of an initial 5'-protected nucleoside is first covalently attached to the polymer support. R.C. Pless *et al.* (1975) *Nucleic Acids Res.* **2**:773 (1975). Synthesis of the oligonucleotide then proceeds by deprotection of the 5'-hydroxyl group of the attached nucleoside, followed by coupling of an incoming nucleoside-3'-phosphoramidite to the deprotected hydroxyl group. M.D. Matteucci *et al.* (1981) *J. Am. Chem. Soc.* **103**:3185. The resulting phosphite triester is finally oxidized to a phosphotriester to complete the internucleotide bond. R.L. Letsinger *et al.* (1976) *J. Am. Chem. Soc.* **98**:3655. The steps of deprotection, coupling and oxidation are repeated until an oligonucleotide of the desired length and sequence is obtained. This process is illustrated schematically in Figure 1 (wherein "B" represents a purine or pyrimidine base, "DMT" represents dimethoxytrityl and "iPR" represents isopropyl).

[0004] The chemical group conventionally used for the protection of nucleoside 5'-hydroxyls is dimethoxytrityl ("DMT"), which is removable with acid. H.G. Khorana (1968) *Pure Appl. Chem.* **17**:349; M. Smith *et al.* (1962) *J. Am. Chem. Soc.* **84**:430. This acid-labile protecting group provides a number of advantages for working with both nucleosides and oligonucleotides. For example, the DMT group can be introduced onto a nucleoside regioselectively and in high yield. E.I. Brown *et al.* (1979) *Methods in Enzymol.* **68**:109. Also, the lipophilicity of the DMT group greatly increases the solubility of nucleosides in organic solvents, and the carbocation resulting from acidic deprotection gives a strong chromophore, which can be used to indirectly monitor coupling efficiency. M.D. Matteucci *et al.* (1980) *Tetrahedron Lett.* **21**:719. In addition, the hydrophobicity of the group can be used to aid separation on reverse-phase HPLC. C. Becker *et al.* (1985) *J. Chromatogr.* **326**:219.

[0005] However, use of DMT as a hydroxyl-protecting group in oligonucleotide synthesis is also problematic. The N-glycosidic linkages of oligodeoxyribonucleotides are susceptible to acid catalyzed cleavage (N.K. Kochetkov *et al.*, *Organic Chemistry of Nucleic Acids* (New York: Plenum Press, 1972)); and even when the protocol is optimized, recurrent removal of the DMT group with acid during oligonucleotide synthesis results in depurination. H. Shaller *et al.* (1963) *J. Am. Chem. Soc.* **85**:3821. The N-6-benzoyl-protected deoxyadenosine nucleotide is especially susceptible to glycosidic cleavage, resulting in a substantially reduced yield of the final oligonucleotide. J.W. Efcavitch *et al.* (1985) *Nucleosides & Nucleotides* **4**:267. Attempts have been made to address the problem of acid-catalyzed depurination utilizing alternative mixtures of acids and various solvents; see, for example, E. Sonveaux (1986) *Bioorganic Chem.* **14**:274. However, this approach has met with limited success. L.J. McBride *et al.* (1986) *J. Am. Chem. Soc.* **108**:2040.

[0006] Conventional synthesis of oligonucleotides using DMT as a protecting group is problematic in other ways as well. For example, cleavage of the DMT group under acidic conditions gives rise to the resonance-stabilized and long-lived bis(p-anisyl)phenylmethyl carbocation. P.T. Gilham *et al.* (1959) *J. Am. Chem. Soc.* **81**:4647. Protection and deprotection of hydroxyl groups with DMT are thus readily reversible reactions, resulting in side reactions during oligonucleotide synthesis and a lower yield than might otherwise be obtained. To circumvent such problems, large excesses of acid are used with DMT to achieve quantitative deprotection. As bed volume of the polymer is increased in larger scale synthesis, increasingly greater quantities of acid are required. The acid-catalyzed depurination which occurs during the synthesis of oligodeoxyribonucleotides is thus increased by the scale of synthesis. M.H. Caruthers *et al.*, in *Genetic Engineering: Principles and Methods*, J.K. Setlow *et al.*, Eds. (New York: Plenum Press, 1982).

[0007] Considerable effort has been directed to developing 5'-O-protecting groups which can be removed under non-acidic conditions. For example, R.L. Letsinger *et al.* (1967) *J. Am. Chem. Soc.* **89**:7147, describe use of a hydrazine-labile benzoyl-propionyl group, and J.F.M. deRoos *et al.* (1979) *Real. Track. Chain. Pays-Bas.* **98**:537, describe using the hydrazine-labile levulinyl ester for 5'-OH protection (see also S. Iwai *et al.* (1988) *Tetrahedron Lett.* **29**:5383; and S. Iwai *et al.* (1988) *Nucleic Acids Res.* **16**:9443). However, the cross-reactivity of hydrazine with pyrimidine nucleotides (as described in F. Baron *et al.* (1955) *J. Chem. Soc.* 2855 and in V. Habermann (1962) *Biochem. Biophys. Acta* **55**:999), the poor selectivity of levulinic anhydride and hydrazine cleavage of N-acyl protecting groups (R.L. Letsinger *et al.* (1968), *Tetrahedron Lett.* **22**:2621) have made these approaches impractical. H. Seliger *et al.* (1985), *Nucleosides & Nucleotides* **4**:153, describes the 5'-O-phenyl-azophenyl carbonyl ("PAPco") group, which is removed by a two-step procedure involving transesterification followed by β -elimination; however, unexpectedly low and non-reproducible yields resulted. Fukuda *et al.* (1988) *Nucleic Acids Res. Symposium Ser.* **19**, 13, and C. Lehmann *et al.* (1989) *Nucleic Acids Res.* **17**:2389, describe application of the 9-fluorenylmethylcarbonate ("Fmoc") group for 5'-protection. C. Leh-

mann *et al.* (1989) report reasonable yields for the synthesis of oligonucleotides up to 20 nucleotides in length. The basic conditions required for complete deprotection of the Fmoc group, however, lead to problems with protecting group compatibility. Similarly, R.L. Letsinger *et al.* (1967), *J. Am. Chem. Soc.* 32:296, describe using the p-nitrophenyloxy-carbonyl group for 5'-hydroxyl protection. In all of the procedures described above utilizing base-labile 5'-O-protecting groups, the requirements of high basicity and long deprotection times have severely limited their application for routine synthesis of oligonucleotides.

[0008] Still an additional drawback associated with conventional oligonucleotide synthesis using DMT as a hydroxyl-protecting group is the necessity of multiple steps, particularly the post-synthetic deprotection step in which the DMT group is removed following oxidation of the internucleotide phosphite triester linkage to a phosphorotriester. It would be desirable to work with a hydroxyl-protecting group that could be removed via oxidation, such that the final two steps involved in nucleotide addition, namely oxidation and deprotection, could be combined.

[0009] The problems associated with the use of DMT are exacerbated in solid phase oligonucleotide synthesis where "microscale" parallel reactions are taking place on a very dense, packed surface. Applications in the field of genomics and high throughput screening have fueled the demand for precise chemistry in such a context. Thus, increasingly stringent demands are placed on the chemical synthesis cycle as it was originally conceived, and the problems associated with conventional methods for synthesizing oligonucleotides are rising to unacceptable levels in these expanded applications.

[0010] The invention is thus addressed to the aforementioned deficiencies in the art, and provides a novel method for synthesizing oligonucleotides, wherein the method has numerous advantages relative to prior methods such as those discussed above. The novel method involves the use of neutral or mildly basic-conditions to remove hydroxyl-protecting groups, such that acid-induced depurination is avoided. In addition, the reagents used provide for irreversible deprotection, significantly reducing the likelihood of unwanted side reactions and increasing the overall yield of the desired product. The method provides for simultaneous oxidation of the internucleoside phosphite triester linkage and removal of the hydroxyl-protecting group, eliminating the extra step present in conventional processes for synthesizing oligonucleotides; the method also avoids the extra step of removing exocyclic amine protecting groups, as the reagents used for hydroxyl group deprotection substantially remove exocyclic amine protecting groups. In addition, the method can be used in connection with fluorescent or other readily detectable protecting groups, enabling monitoring of individual reaction steps. Further, the method is useful in carrying out either 3'-to-5' synthesis or 5'-to-3' synthesis. Finally, because of the far more precise chemistry enabled by the present invention, the method readily lends itself to the highly parallel, microscale synthesis of oligonucleotides.

[0011] It is accordingly a primary object of the invention to provide a novel method for synthesizing oligonucleotides which addresses and overcomes the above-mentioned disadvantages of the prior art.

[0012] It is another object of the invention to provide a novel method for synthesizing oligonucleotides in which individual nucleoside monomers are added to a growing oligonucleotide chain using carbonates as hydroxyl-protecting groups and "alpha effect" nucleophiles as deprotecting reagents.

[0013] It is still another object of the invention to provide such a method in which hydroxyl group deprotection and oxidation of the internucleotide phosphite triester linkage are carried out simultaneously, in a single step.

[0014] It is yet another object of the invention to provide such a method in which deprotection and oxidation are conducted in aqueous solution at neutral or mildly basic pH.

[0015] It is an additional object of the invention to provide such a method in which removal of hydroxyl protecting groups during oligonucleotide synthesis is irreversible.

[0016] It is a further object of the invention to provide such a method in which the desired oligonucleotide can be synthesized in either the 3'-to-5' direction or the 5'-to-3' direction.

[0017] Still a further object of the invention is to provide such a method in which individual oligonucleotides are synthesized within the context of a highly dense, substantially parallel oligonucleotide array on a substrate surface.

[0018] Still an additional object of the invention is to provide nucleoside reagents useful in conjunction with the novel synthetic methods.

[0019] Additional objects, advantages and novel features of the invention will be set forth in part in the description which follows, and in part will become apparent to those skilled in the art upon examination of the following, or may be learned by practice of the invention.

[0020] The invention is premised on the discovery that rapid and selective removal of suitable 5'-OH or 3'-OH protecting groups following phosphoramidite condensation can be achieved by employing nucleophiles, and particularly peroxy anions, that exhibit an "alpha effect" under neutral or mildly basic conditions. Further, it has now been discovered that rapid and selective deprotection can be achieved under such conditions by employing carbonate groups for 5'-OH or 3'-OH protection. Deprotection of nucleoside carbonates using peroxy anions can be conducted in aqueous solution, at neutral or mild pH, resulting in quantitative removal of the carbonate group and concomitant and quantitative oxidation of the internucleotide phosphite triester bond. Oligonucleotides synthesized using the novel methodology can be isolated in high yield and substantially free of detectable nucleoside modifications.

[0021] The term "alpha effect," as in an "alpha effect" nucleophilic deprotection reagent, is used to refer to an enhancement of nucleophilicity that is found when the atom adjacent a nucleophilic site bears a lone pair of electrons. As the term is used herein, a nucleophile is said to exhibit an "alpha effect" if it displays a positive deviation from a Brønsted-type nucleophilicity plot. S. Hoz *et al.* (1985) *Israel J Chem.* **26**:313. See also, J.D. Aubort *et al.* (1970) *Chem. Comm.* 1378; J.M. Brown *et al.* (1979) *J. Chem. Soc. Chem. Comm.* 171; E. Buncl *et al.* (1982) *J. Am. Chem. Soc.* **104**:4896; J.O. Edwards *et al.* (1962) *J. Amer. Chem. Soc.* **84**:16; J.D. Evanseck *et al.* (1987) *J. Am. Chem. Soc.* **109**:2349. The magnitude of the alpha effect is dependent upon the electrophile which is paired with the specific nucleophile. J.E. McIsaac, Jr. *et al.* (1972), *J. Org. Chem.* **37**:1037. Peroxy anions are example of nucleophiles which exhibit strong alpha effects.

[0022] In one general-aspect, the invention features a method, in an oligonucleotide synthesis, for removing a protecting group from a protected nucleoside, by reacting the protected nucleoside or protected nucleotide with a nucleophile that exhibits an alpha effect at conditions of mildly basic pH, and particularly at conditions of pH 10 or less.

[0023] The invention provides for efficient solid-phase synthesis of oligonucleotides of lengths up to 25 nucleotides and greater. Treatment using an alpha effect nucleophile according to the invention for removal of carbonate protecting groups is irreversible, resulting in breakdown of the carbonate and formation of CO₂. Moreover, because such treatment results in concomitant oxidation of the internucleotide bond and substantial removal of exocyclic amine protecting groups, the method of the invention obviates the need for a separate oxidation step and a post-synthesis deprotection step to remove any exocyclic amine protecting groups that may be used.

[0024] In another general aspect, the invention features a method for making an oligonucleotide array made up of array features each presenting a specified oligonucleotide sequence at an address on an array substrate, by first treating the array substrate to protect the hydroxyl moieties on the derivatized surface from reaction with phosphoramidites, then carrying out the steps of (a) applying droplets of an alpha effect nucleophile to effect deprotection of hydroxyl moieties at selected addresses, and (b) flooding the array substrate with a medium containing a selected protected phosphoramidite to permit coupling of the selected phosphoramidite onto the deprotected hydroxyl moieties at the selected addresses, and repeating the steps (a) and (b) to initiate and to sequentially build up oligonucleotides having the desired sequences at the selected addresses to complete the array features. In a variation on the aforementioned method, the droplets applied may comprise the protected phosphoramidite, and the alpha effect nucleophile may be used to flood the array substrate.

[0025] In the array construction method according to the invention, the deprotection reagents are aqueous, allowing for good droplet formation on a wide variety of array substrate surfaces. Moreover, because the selection of features employs aqueous media, small-scale discrete droplet application onto specified array addresses can be carried out by adaptation of techniques for reproducible fine droplet deposition from printing technologies. Further, as noted above, the synthesis reaction provides irreversible deprotection resulting in evolution of CO₂, and thus quantitative removal of protecting groups within each droplet may be achieved. The phosphoramidite reactions are carried out in bulk, employing an excess of the phosphoramidite during the coupling step (b), allowing for exclusion of water by action of the excess phosphoramidite as a desiccant.

[0026] In a first aspect of the invention, there is provided a method for synthesizing oligonucleotides which comprises (a) condensing the 3' -OH or 5' -OH group of a support-bound nucleoside or oligonucleotide with a monomeric nucleoside phosphoramidite having a carbonate-protected hydroxyl group, to provide an intermediate in which the support-bound nucleoside or oligonucleotide is bound to the monomeric nucleoside through a phosphite triester linkage, and (b) treating the intermediate provided in step (a) with a deprotecting reagent effective to convert the carbonate-protected hydroxyl group to a free hydroxyl moiety and simultaneously oxidize the phosphite triester linkage to give a phosphotriester linkage.

[0027] Preferably, (a) and (b) are carried out in aqueous solution.


[0028] Preferably, step (b) is an irreversible reaction.

[0029] Preferably, step (b) is conducted at neutral to mildly basic pH, in which case it is preferred that the deprotecting reagent used in step (b) comprises a nucleophile that exhibits an alpha effect at neutral to mildly basic pH. Step (b) may be conducted at a pH of less than about 10.


[0030] In a preferred embodiment, the nucleophile is a peroxide, and preferably the peroxide is an inorganic peroxide of the formula M⁺OOH⁻, wherein M⁺ is a counterion. The counterion may be selected from the group consisting of H⁺, Li⁺, Na⁺, K⁺, Rb⁺ and Cs⁺.

[0031] In one embodiment, the peroxide is an organic peroxide of the formula ROOH, wherein R is hydrocarbyl optionally substituted with one or more nonhydrocarbyl substituents and optionally containing one or more nonhydrocarbyl linkages. The peroxide may have the structure of formula (V), (VI) or (VII)

(V)



(VI)

(VII) $R^{10}-O-O^-$

in which R^4 through R^{10} are hydrocarbyl optionally substituted with one or more nonhydrocarbyl substituents and optionally containing one or more nonhydrocarbyl linkages.

[0032] In an alternative embodiment, the peroxide is one of t-butyl hydroperoxide, m-chloroperoxybenzoic acid, and mixtures thereof.


[0033] Preferably, step (b) is conducted at a pH at which said nucleophile exhibits a maximum alpha effect, and preferably is conducted at a pH that is greater than the pKa of said nucleophile.

[0034] Steps (a) and (b) may be repeated until an oligonucleotide having a desired sequence and length is provided.

[0035] In one embodiment, synthesis is conducted in the 3'-to-5' direction, and step (a) involves condensation at the 5'-OH of the support-bound nucleoside or oligonucleotide.

[0036] The monomeric nucleoside phosphoramidite may have the structure of formula (II)

(II)



wherein:


B is a purine or pyrimidine base;

R is hydrido or hydroxyl;

R^1 is $COOR^3$, wherein R^3 is substituted or unsubstituted hydrocarbyl;

R^2 has the structure (III)

(III)



wherein X is NQ^1Q^2 in which Q^1 and Q^2 are independently selected from the group consisting of alkyl, aryl, aralkyl, alkaryl, cycloalkyl, alkenyl, cycloalkenyl, alkynyl or cycloalkynyl, optionally containing one or more nonhydrocarbyl linkages and optionally substituted on one or more available carbon atoms with a nonhydrocarbyl substituent, or wherein Q^1 and Q^2 are linked to form a mono- or polyheterocyclic ring, and Y is hydrido or hydrocarbyl.

[0037] Preferably, X is di(lower alkyl)amino or morpholino and Y is selected from the group consisting of lower alkyl, electron-withdrawing β -substituted aliphatic, electron-withdrawing substituted phenyl, and electron-withdrawing substituted phenylethyl.


[0038] More preferably, Y is selected from the group consisting of methyl, 13-cyanoethyl and 4-nitrophenylethyl.

[0039] Preferably, R¹ is selected from the group consisting of *o*-nitrophenylcarbonyl, *p*-phenylazophenylcarbonyl, phenylcarbonyl, *p*-chlorophenylcarbonyl, 5'-(α -methyl-2-nitropiperonyl)oxycarbonyl ("MeNPOC"), 9-fluorenylmethylcarbonyl ("Fmoc"), 2,2,2-trichloro-1,1-dimethylcarbonyl ("TCBOC") and cyano-substituted lower alkyl carbonates.

[0040] In a particularly preferred embodiment, R³ comprises a moiety that becomes fluorescent or coloured upon cleavage of the carbonate -OCOOR³.

[0041] In an alternative embodiment, synthesis is conducted in the 5'-to -3' direction, and step (a) involves condensation at the 3'-OH of the support-bound nucleoside or oligonucleotide.

[0042] In this embodiment, the monomeric nucleoside phosphoramidite preferably has the structure of formula (X)



wherein:


B is a purine or pyrimidine base;

R is hydrido or hydroxyl;

R¹ is COOR³, wherein R³ is substituted or unsubstituted hydrocarbyl;

R² has the structure (III)

(III)



wherein X is NQ¹Q² in which Q¹ and Q² are independently selected from the group consisting of alkyl, aryl, aralkyl, alkaryl, cycloalkyl, alkenyl, cycloalkenyl, alkenyl or cycloalkenyl, optionally containing one or more nonhydrocarbyl linkages and optionally substituted on one or more available carbon atoms with a nonhydrocarbyl substituent, or wherein Q¹ and Q² are linked to form a mono- or polyheterocyclic ring, and Y is hydrido or hydrocarbyl.

[0043] X is preferably di(lower alkyl)amino or morpholino and Y is selected from the group consisting of lower alkyl, electron-withdrawing β -substituted aliphatic, electron-withdrawing substituted phenyl, and electron-withdrawing substituted phenylethyl.

[0044] More preferably, Y is selected from the group consisting of methyl, β -cyanoethyl and 4-nitrophenylethyl.

[0045] R¹ is preferably selected from the group consisting of *o*-nitrophenylcarbonyl, *p*-phenylazophenylcarbonyl, phenylcarbonyl, *p*-chlorophenylcarbonyl, 5'-(α -methyl-2-nitropiperonyl)oxycarbonyl ("MeNPOC"), 9-fluorenylmethylcarbonyl ("Fmoc"), 2,2,2-trichloro-1,1-dimethylcarbonyl ("TCBOC") and cyano-substituted lower alkyl carbonates.

[0046] R³ preferably comprises a moiety that becomes fluorescent or coloured upon cleavage of the carbonate -OCOOR³.

[0047] In another aspect of the invention, there is provided a method for making an oligonucleotide array made up of array features each presenting a specified oligonucleotide sequence at an address on an array substrate, the method comprising steps of:

providing a hydroxyl-derivatized array substrate and treating the array substrate to protect hydroxyl moieties on the derivatized surface from reaction with phosphoramidites,

then iteratively carrying out the steps of (i) applying droplets of an alpha effect nucleophile to effect deprotection

of hydroxyl moieties at selected addresses, and (ii) flooding the array substrate with a medium containing a selected monomeric nucleoside phosphoramidite having a carbonate-protected hydroxyl group, to permit covalent attachment of the selected nucleoside to the deprotected hydroxyl moieties at the selected addresses.

[0048] Steps (i) and (ii) are preferably carried out in aqueous solution.

[0049] Preferably, step (i) is an irreversible reaction.

Step (i) may be conducted at neutral to mildly basic pH;


[0050] Step (b) is preferably conducted at a pH of less than about 10.

[0051] The nucleophile is preferably a peroxide, and more preferably an inorganic peroxide of the formula M^+OOH^- , wherein M^+ a counterion. The counterion may be selected from the group consisting of H^+ , Li^+ , Na^+ , K^+ , Rb^+ and Cs^+ .


[0052] The peroxide may be an organic peroxide of the formula $ROOH$, wherein R is hydrocarbyl optionally substituted with one or more nonhydrocarbyl substituents and optionally containing one or more nonhydrocarbyl linkages.

[0053] Preferably, the peroxide has the structure of formula (V), (VI) or (VII)

(V)



(VI)




(VII) $R^{10}-O-O^-$

in which R^4 through R^{10} are hydrocarbyl optionally substituted with one or more nonhydrocarbyl linkages.


[0054] In an alternative embodiment, the peroxide is one of t-butyl hydroperoxide, m-chloroperoxybenzoic acid, and mixtures thereof.

[0055] In a further aspect of the inventions there is provided a nucleoside monomer having the structure of formula (II) or formula (X)

(II)



(X)



wherein:


B is a purine pyrimidine base, as described previously herein;

R is hydrido or hydroxyl;

R¹ is COOR³, wherein R³ is substituted or unsubstituted hydrocarbyl; and

R² has the structure (III)

(III)



wherein X is NQ¹Q² in which Q¹ and Q² are independently selected from the group consisting of alkyl, aryl, aralkyl, alkaryl, cycloalkyl, alkenyl, cycloalkenyl, alkenyl or cycloalkynyl, optionally containing one or more nonhydrocarbyl linkages and optionally substituted on one or more available carbon atoms with a nonhydrocarbyl substituent, or wherein Q¹ and Q² are linked to form a mono- or polyheterocyclic ring, and Y is hydrido or hydrocarbyl.

[0056] Preferably, X is di(lower alkyl)amino or morpholino and Y is selected from the group consisting of lower alkyl, electron-withdrawing β -substituted aliphatic, electron-withdrawing substituted phenyl, and electron-withdrawing substituted phenylethyl. More preferably, Y is selected from the group consisting of methyl 13-cyanoethyl and 4-nitrophenylethyl.

[0057] R³ is preferably selected from the group consisting of *o*-nitrophenylcarbonyl, *p*-phenylazophenylcarbonyl, phenylcarbonyl, *p*-chlorophenylcarbonyl, 5'-(α -methyl-2-nitropiperonyl)oxycarbonyl ("MeNPOC"), 9-fluorenylmethylcarbonyl ("Fmoc"), 2,2,2-trichloro-1,1-dimethylcarbonyl ("TCBOC") and cyano-substituted lower alkyl carbonates.

[0058] In a preferred embodiment, R³ comprises a moiety that becomes fluorescent or coloured upon cleavage of the carbonate -OCOOR³.

[0059] A particularly preferred nucleoside monomer is that having the structure of formula (II) or the structure of formula (X).

[0060] In yet a further aspect of the invention, there is provided a kit for synthesizing an oligonucleotide on a solid support, comprising a hydroxyl-derivatized support surface, a protecting group for protecting hydroxyl moieties on said derivatized support surface, at least one protected nucleoside, at least one nucleoside phosphoramidite, a nucleophile that exhibits an alpha effect at neutral to mildly basic pH, and reagents suitable for establishing pH and for carrying out reactions of deprotection, phosphoramidite coupling, and oxidation to form an internucleotide phosphotriester linkage.

[0061] Figure 1 schematically illustrates conventional 3'-to-5' oligonucleotide synthesis using DMT as a 5'-OH protecting group, and separate deprotection and oxidation steps.

[0062] Figure 2 schematically illustrates 3'-to-5' oligonucleotide synthesis using the method of the invention.

[0063] Figures 3A and 3B compare the conventional deprotection reaction in which DMT is used as a hydroxyl-protecting group (Figure 3A) and the deprotection reaction in which the reagents of the invention are employed (Figure 3B).

[0064] Figure 4 schematically illustrates a method for synthesizing a 5'-carbonate-3'-phosphoramidite monomer of the invention.

[0065] Figure 5 schematically illustrates a method for synthesizing a 3'-carbonate-5'-phosphoramidite monomer of the invention.

[0066] Figure 6 sets forth the HPLC results obtained for a mixed-sequence oligonucleotide synthesized in Example 4, part (D).

[0067] Figure 7 sets forth the MALDI TOF results obtained for the same mixed-sequence oligonucleotide.

[0068] It is to be understood that unless otherwise indicated, this invention is not limited to specific reagents, reaction conditions, synthetic steps, or the like, as such may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only and is not intended to be limiting.

[0069] It must be noted that, as used in the specification and the appended claims, the singular forms "a," "an" and "the" include plural referents unless the context clearly dictates otherwise. Thus, for example, reference to "a protecting group" includes combinations of protecting groups, reference to "a nucleoside" includes combinations of nucleosides, and the like. Similarly, reference to "a substituent" as in a compound substituted with "a substituent" includes the possibility of substitution with more than one substituent, wherein the substituents may be the same or different.

[0070] In this specification and in the claims which follow, reference will be made to a number of terms which shall be defined to have the following meanings:

[0071] The term "alkyl" as used herein, unless otherwise specified, refers to a saturated straight chain, branched or cyclic hydrocarbon group of 1 to 24, typically 1-12, carbon atoms, such as methyl, ethyl, *n*-propyl, isopropyl, *n*-butyl, isobutyl, *t*-butyl, pentyl, cyclopentyl, isopentyl, neopentyl, hexyl, isohexyl, cyclohexyl, 3-methylpentyl, 2,2-dimethylbutyl, and 2,3-dimethylbutyl. The term "lower alkyl" intends an alkyl group of one to six carbon atoms, and includes, for example, methyl, ethyl, *n*-propyl, isopropyl, *n*-butyl, isobutyl, *t*-butyl, pentyl, cyclopentyl, isopentyl, neopentyl, hexyl, isohexyl, cyclohexyl, 3-methylpentyl, 2,2-dimethylbutyl, and 2,3-dimethylbutyl. The term "cycloalkyl" refers to cyclic alkyl groups such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl and cyclooctyl.

[0072] The term "alkenyl" as used herein, unless otherwise specified, refers to a branched, unbranched or cyclic (in the case of C₅ and C₆) hydrocarbon group of 2 to 24, typically 2 to 12, carbon atoms containing at least one double bond, such as ethenyl, vinyl, allyl, octenyl, decenyl, and the like. The term "lower alkenyl" intends an alkenyl group of two to six carbon atoms, and specifically includes vinyl and allyl. The term "cycloalkenyl" refers to cyclic alkenyl groups.

[0073] The term "alkynyl" as used herein, unless otherwise specified, refers to a branched or unbranched hydrocarbon group of 2 to 24, typically 2 to 12, carbon atoms containing at least one triple bond, such as acetylenyl, ethynyl, *n*-propynyl, isopropynyl, *n*-butynyl, isobutynyl, *t*-butynyl, octynyl, decynyl and the like. The term "lower alkynyl" intends an alkynyl group of two to six carbon atoms, and includes, for example, acetylenyl and propynyl, and the term "cycloalkynyl" refers to cyclic alkynyl groups.

[0074] The term "aryl" as used herein refers to an aromatic species containing 1 to 5 aromatic rings, either fused or linked, and either unsubstituted or substituted with 1 or more substituents typically selected from the group consisting of amino, halogen and lower alkyl. Preferred aryl substituents contain 1 to 3 fused aromatic rings, and particularly preferred aryl substituents contain 1 aromatic ring or 2 fused aromatic rings. Aromatic groups herein may or may not be heterocyclic. The term "aralkyl" intends a moiety containing both alkyl and aryl species, typically containing less than about 24 carbon atoms, and more typically less than about 12 carbon atoms in the alkyl segment of the moiety, and typically containing 1 to 5 aromatic rings. The term "aralkyl" will usually be used to refer to aryl-substituted alkyl groups. The term "aralkylene" will be used in a similar manner to refer to moieties containing both alkylene and aryl species, typically containing less than about 24 carbon atoms in the alkylene portion and 1 to 5 aromatic rings in the aryl portion, and typically aryl-substituted alkylene. Exemplary aralkyl groups have the structure -(CH₂)_j-Ar wherein j is an integer in the range of 1 to 24, more typically 1 to 6, and Ar is a monocyclic aryl moiety.

[0075] The term "electron withdrawing" denotes the tendency of a substituent to attract valence electrons of the molecule of which it is a part, i.e., an electron-withdrawing substituent is electronegative.

[0076] The term "heterocyclic" refers to a five- or six-membered monocyclic structure or to an eight- to eleven-membered bicyclic structure which is either saturated or unsaturated. The heterocyclic groups herein may be aliphatic or aromatic. Each heterocycle consists of carbon atoms and from one to four heteroatoms selected from the group consisting of nitrogen, oxygen and sulfur. As used herein, the terms "nitrogen heteroatoms" and "sulfur heteroatoms" include any oxidized form of nitrogen and sulfur, and the quaternized form of any basic nitrogen. Examples of heterocyclic groups include piperidinyl, morpholinyl and pyrrolidinyl.

[0077] The term "halo" or "halogen" is used in its conventional sense to refer to a chloro, bromo, fluoro or iodo substituent.

[0078] As used herein, the term "oligonucleotide" shall be generic to polydeoxynucleotides (containing 2-deoxy-D-ribose), to polyribonucleotides (containing D-ribose), to any other type of polynucleotide which is an N-glycoside of a purine or pyrimidine base, and to other polymers containing nonnucleotidic backbones, providing that the polymers contain nucleobases in a configuration which allows for base pairing and base stacking, such as is found in DNA and RNA.

[0079] It will be appreciated that, as used herein, the terms "nucleoside" and "nucleotide" will include those moieties which contain not only the known purine and pyrimidine bases, but also modified purine and pyrimidine bases and other heterocyclic bases which have been modified (these moieties are sometimes referred to herein, collectively, as "purine and pyrimidine bases and analogs thereof"). Such modifications include methylated purines or pyrimidines, acylated purines or pyrimidines, and the like.

[0080] By "protecting group" as used herein is meant a species which prevents a segment of a molecule from undergoing a specific chemical reaction, but which is removable from the molecule following completion of that reaction. This is in contrast to a "capping group," which permanently binds to a segment of a molecule to prevent any further chemical transformation of that segment.

[0081] "Optional" or "optionally" means that the subsequently described circumstance may or may not occur, so that the description includes instances where the circumstance occurs and instances where it does not. For example, the phrase "optionally substituted" means that a non-hydrogen substituent may or may not be present, and, thus, the description includes structures wherein a non-hydrogen substituent is present and structures wherein a non-hydrogen substituent is not present.

[0082] In a first embodiment, the invention pertains to a method for synthesizing an oligonucleotide on a solid support, wherein a carbonate is used as a hydroxyl-protecting group and an alpha effect nucleophile is used to bring about deprotection. The novel synthesis is based on a simple, two-step method of (1) coupling a hydroxyl-protected nucleoside monomer to a growing oligonucleotide chain, and (2) deprotecting the product, under neutral or mildly basic conditions, using an alpha effect nucleophilic reagent that also oxidizes the internucleotide linkage to give a phosphotriester bond. The coupling and deprotection/oxidation steps are repeated as necessary to give an oligonucleotide having a desired sequence and length.

[0083] In the initial step of the synthesis, then, an unprotected nucleoside is covalently attached to a solid support to serve as the starting point for oligonucleotide synthesis. The nucleoside may be bound to the support through its 3'-hydroxyl group or its 5'-hydroxyl group, but is typically bound through the 3'-hydroxyl group. A second nucleoside monomer is then coupled to the free hydroxyl group of the support-bound initial monomer, wherein for 3'-to-5' oligonucleotide synthesis, the second nucleoside monomer has a phosphorus derivative such as a phosphoramidite at the 3' position and a carbonate protecting group at the 5' position, and alternatively, for 5'-to-3' oligonucleotide synthesis, the second nucleoside monomer has a phosphorus derivative at the 5' position and a carbonate protecting group at the 3' position. This coupling reaction gives rise to a newly formed phosphite triester bond between the initial nucleoside monomer and the added monomer, with the carbonate-protected hydroxyl group intact. In the second step of the synthesis, the carbonate group is removed with an alpha effect nucleophile that also serves to oxidize the phosphite triester linkage to the desired phosphotriester.

[0084] More specifically, for 3'-to-5' synthesis, a support-bound nucleoside monomer is provided having the structure (I)



wherein:

O represents the solid support or a support-bound oligonucleotide chain;

R is hydrido or hydroxyl, wherein when R is hydrido, the support-bound nucleoside is a deoxyribonucleoside, as will be present in DNA synthesis, and when R is hydroxyl, the support-bound nucleoside is a ribonucleoside, as will be present in RNA synthesis; and


B is a purine or pyrimidine base. The purine or pyrimidine base may be conventional, e.g., adenine (A), thymine (T), cytosine (C), guanine (G) or uracil (U), or a protected form thereof, e.g., wherein the base is protected with a protecting group such as acetyl, difluoroacetyl, trifluoroacetyl, isobutyryl, benzoyl, or the like. The purine or pyrimidine base may also be an analog of the foregoing; suitable analogs will be known to those skilled in the art and are described in the pertinent texts and literature. Common analogs include, but are not limited to, 1-methyladenine, 2-methyladenine, N⁶-methyladenine, N⁶-isopentyladenine, 2-methylthio-N⁶-isopentyladenine, N,N-dimethyladenine, 8-bromoadenine, 2-thiocytosine, 3-methylcytosine, 5-methylcytosine, 5-ethylcytosine, 4-acetylcytosine, 1-methylguanine, 2-methylguanine, 7-methylguanine, 2,2-dimethylguanine, 8-bromoguanine, 8-chloroguanine, 8-aminoguanine, 8-methylguanine, 8-thioguanine, 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, 5-ethyluracil, 5-propyluracil, 5-methoxyuracil, 5-hydroxymethyluracil, 5-(carboxyhydroxymethyl)uracil, 5-(methylaminomethyl)uracil, 5-(carboxymethylaminomethyl)uracil, 2-thiouracil, 5-methyl-2-thiouracil, 5-(2-bromovinyl)

uracil, uracil-5-oxyacetic acid, uracil-5-oxyacetic acid methyl ester, pseudouracil, 1-methylpseudouracil, queosine, inosine, 1-methylinosine, hypoxanthine, xanthine, 2-aminopurine, 6-hydroxyaminopurine, 6-thiopurine and 2,6-diaminopurine.

[0085] The protected monomer to be added has the structure of formula (II)




in which B and R are as defined above with respect to the support-bound nucleoside of structural formula (I), and R¹ is COOR³, such that a carbonate group -OCOOR³ is present at the 5' position. R³ is generally substituted or unsubstituted hydrocarbyl, including alkyl, aryl, aralkyl, alkaryl, cycloalkyl, alkenyl, cycloalkenyl, alkynyl, cycloalkynyl, optionally containing one or more nonhydrocarbyl linkages such as ether linkages, thioether linkages, oxo linkages, amine and imine linkages, and optionally substituted on one or more available carbon atoms with a nonhydrocarbyl substituent such as cyano, nitro, halo, or the like. Preferred carbonate groups -OCOOR³ are aryl carbonates, i.e., R³ is aryl. Suitable aryl carbonates include, for example, *o*-nitrophenylcarbonyl, *p*-phenylazophenylcarbonyl, phenylcarbonyl, *p*-chlorophenylcarbonyl, 5'-(α -methyl-2-nitropiperonyl)oxycarbonyl ("MeNPOC"), and 9-fluorenylmethylcarbonyl ("Fmoc"). Particularly preferred aryl carbonates have the structure Ar-L-O-(CO)-O- wherein Ar is an aromatic moiety; typically a monocyclic aromatic moiety such as a phenyl group, optionally substituted with one or more electron-withdrawing substituents such as halo, nitro, cyano, or the like, and L is a lower alkylene linkage. Preferred alkyl carbonate substituents are fluorinated alkyl carbonates such as 2,2,2-trichloro-1,1-dimethylcarbonyl ("TCBOC") and cyano-substituted alkyl carbonates such as 1,1-dimethyl-2-cyanoethyl carbonate



[0086] R³ may also be a fluorescent or colored moiety. Preferably, in this embodiment, R³ becomes fluorescent or colored upon cleavage of the carbonate -OCOOR³, but is neither fluorescent nor colored when bound to the nucleoside in carbonate form. In this way, when the carbonate protecting group R¹ is removed, the reaction may be monitored by detecting a fluorescent or colored cleavage product. Alternatively, R³ may be fluorescent or colored when bound to the nucleoside in carbonate form, such that the presence of the newly attached monomer can be immediately detected. Examples of fluorescent and colorimetric species that may be employed include, but are not limited to: xanthenes such as fluoresceins, eosins and erythrosins, with preferred fluorescein compounds exemplified by 6-carboxy-fluorescein, 5- and 6-carboxy-4,7-dichlorofluorescein, 2',7'-dimethoxy-5- and 6-carboxy-4,7-dichlorofluorescein, 2',7'-dimethoxy-4',5'-dichloro-5- and 6-carboxyfluorescein, 2',7'-dimethoxy-4',5'-dichloro-5- and 6-carboxy-4,7-dichlorofluorescein, 1', 2',7',8'-dibenzo-5- and 6-carboxy-4,7-dichlorofluorescein, 2',7'-dichloro-5- and 6-carboxy-4,7-dichlorofluorescein, and 2',4',5',7'-tetrachloro-5- and 6-carboxy-4,7-dichlorofluorescein; rhodamines such as tetramethylrhodamine and Texas Red®; benzimidazoles; ethidium; propidium; anthracyclines; mithramycins; acridines; actinomycins; merocyanines; coumarins such as 4-methyl-7-methoxycoumarin; pyrenes; chrysenes; stilbenes; anthracenes; naphthalenes such as dansyl, 5-dimethylamino-1-naphthalenesulfonyl; salicylic acids; benz-2-oxa-1-diazoles (also known as benzofurans), including 4-amino-7-nitrobenz-2-oxa-1,3-diazole; fluorescamine; and 4-methylumbelliferone.

[0087] R² is a phosphorus derivative that enables coupling to a free hydroxyl group. Preferred phosphorus derivatives are phosphoramidites, such that R² has the structure (III)

(III)




wherein X is NQ^1Q^2 in which Q^1 and Q^2 may be the same or different and are typically selected from the group consisting of alkyl, aryl, aralkyl, alkaryl, cycloalkyl, alkenyl, cycloalkenyl, alkynyl, cycloalkynyl, optionally containing one or more nonhydrocarbyl linkages such as ether linkages, thioether linkages, oxo linkages, amine and imine linkages, and optionally substituted on one or more available carbon atoms with a nonhydrocarbyl substituent such as cyano, nitro, halo, or the like. Preferably, Q^1 and Q^2 represent lower alkyl, more preferably sterically hindered lower alkyls such as isopropyl, *t*-butyl, isobutyl, *sec*-butyl, neopentyl, *tert*-pentyl, isopentyl, *sec*-pentyl, and the like. Most preferably, Q^1 and Q^2 both represent isopropyl. Alternatively, Q^1 and Q^2 may be linked to form a mono- or polyheterocyclic ring having a total of from 1 to 3, usually 1 to 2 heteroatoms and from 1 to 3 rings. In such a case, Q^1 and Q^2 together with the nitrogen atom to which they are attached represent, for example, pyrrolidone, morpholino or piperidino. Usually, Q^1 and Q^2 have a total of from 2 to 12 carbon atoms. Examples of specific $\text{-NQ}^1\text{Q}^2$ moieties thus include, but are not limited to, dimethylamine, diethylamine, diisopropylamine, dibutylamine, methylpropylamine, methylhexylamine, methylcyclopropylamine, ethylcyclohexylamine, methylbenzylamine, methylcyclohexylmethylamine, butylcyclohexylamine, morpholine, thiomorpholine, pyrrolidine, piperidine, 2,6-dimethylpiperidine, piperazine, and the like.


[0088] The moiety "Y" is hydrido or hydrocarbyl, typically alkyl, alkenyl, aryl, aralkyl, or cycloalkyl. Preferably, Y represents: lower alkyl; electron-withdrawing β -substituted aliphatic, particularly electron-withdrawing β -substituted ethyl such as β -trihalomethyl ethyl, β -cyanoethyl, β -sulfoethyl, β -nitro-substituted ethyl, and the like; electron-withdrawing substituted phenyl, particularly halo-, sulfo-, cyano- or nitro-substituted phenyl; or electron-withdrawing substituted phenylethyl. Most preferably, Y represents methyl, β -cyanoethyl, or 4-nitrophenylethyl.

[0089] The coupling reaction is conducted under standard conditions used for the synthesis of oligonucleotides and conventionally employed with automated oligonucleotide synthesizers. Such methodology will be known to those skilled in the art and is described in the pertinent texts and literature, e.g., in D.M. Matteuci et al. (1980) *Tet. Lett.* 521:719 and U.S. Patent No. 4,500,707. The product of the coupling reaction may be represented as structural formula (IV), as follows:

(IV)




[0090] In the second step of the synthesis, the product (IV) is treated with an "alpha effect" nucleophile in order to remove the carbonate protecting group at the 5' terminus, thus converting the moiety -OR^1 to -OH . The alpha effect nucleophile also oxidizes the newly formed phosphite triester linkage -O-P(OY)-O- to give the desired phosphotriester linkage




[0091] Advantageously, this step is conducted in an aqueous solution at neutral pH or at a mildly basic pH, depending on the pKa of the nucleophilic deprotection reagent. That is, and as will be explained in further detail below, the pH at which the deprotection reaction is conducted must be above the pKa of the deprotection reagent for the reagent to be effective. Typically, the reaction is conducted at a pH of less than about 10.

[0092] In a preferred embodiment, the nucleophilic deprotection reagent that exhibits an alpha effect is a peroxide or a mixture of peroxides, and the pH at which deprotection is conducted is at or above the pKa for formation of the corresponding peroxy anion. The peroxide may be either inorganic or organic. Suitable inorganic peroxides include those of the formula M^+OOH^- , where M is any counteranion, including for example H^+ , Li^+ , Na^+ , K^+ , Rb^+ , Cs^+ , or the like; and lithium peroxide or hydrogen peroxide can be particularly suitable. Suitable organic peroxides include those of the formula ROOH , where R is selected from the group consisting of alkyl, aryl, substituted alkyl and substituted aryl. More particularly, the organic peroxide will have one of the following three general structures (V), (VI) or (VII)

(V)




(VI)

(VII) $\text{R}^{10} \text{---} \text{O} \text{---} \text{O}^-$


in which R^4 through R^{10} are generally hydrocarbyl optionally substituted with one or more nonhydrocarbyl substituents and optionally containing one or more nonhydrocarbyl linkages. Generally, R^4 through R^{10} are independently selected from the group consisting of hydrido, alkyl, aryl, aralkyl, cycloalkyl, cycloalkylalkyl, alkenyl, cycloalkenyl, alkynyl, aralkynyl, cycloalkynyl, substituted aralkyl, substituted cycloalkyl, substituted cycloalkylalkyl, substituted alkenyl, substituted cycloalkenyl, substituted alkynyl, substituted aralkynyl, substituted cycloalkynyl; *t*-butyl-hydroperoxide or metachloroperoxybenzoic acid can be particularly suitable. As a specific example, the *m*-chloroperoxybenzoic acid (mCPBA) peroxy anion exhibits a strong alpha effect towards the *p*-chlorophenylcarbonate electrophile, and that, accordingly, the peroxy anion of mCPBA is a particularly effective deprotection reagent for removal of *p*-chlorophenylcarbonate protecting groups.

[0093] The product of this simultaneous deprotection and oxidation step may thus be represented as follows:

(VIII)



wherein B, R and Y are as defined earlier herein. This latter reaction also gives rise to the byproducts R^3O^- and carbon dioxide, insofar as nucleophilic attack of the peroxide deprotection reagent cleaves the carbonate linkage as follows:



[0094] The use of a peroxy anion to effect simultaneous removal of the carbonate protecting group and oxidation of the internucleotide linkage also removes, to a large extent, exocyclic amine-protecting groups such as acetyl, trifluoroacetyl, difluoroacetyl and trifluoroacetyl moieties. Thus, an added advantage herein is the elimination of a separate post-synthetic reaction step to remove exocyclic amine-protecting groups, as is required with conventional methods of synthesizing oligonucleotides. Elimination of this additional step significantly decreases the time and complexity involved in oligonucleotide synthesis.


[0095] An additional advantage of peroxy anions as deprotection reagents herein is that they may be readily activated or inactivated by simply changing pH. That is, the effectiveness of peroxides as nucleophiles is determined by their pKa. In buffered solutions having a pH below the pKa of a particular peroxide, the peroxides are not ionized and thus are non-nucleophilic. To activate a peroxide and render it useful as a deprotection reagent for use herein, the pH is increased above the pKa so that the peroxide is converted to a nucleophilic peroxy anion. Thus, one can carefully control the timing and extent of the deprotection reaction by varying the pH of the peroxide solution used.

[0096] Figure 2 schematically illustrates 3'-to-5' synthesis of an oligonucleotide using the method of the present invention. In the figure, the moiety "Arco" ("aryloxycarbonyl") represents the carbonate protecting group *p*-chlorophenylcarbonyl. As may be seen, deprotection and oxidation occur simultaneously. The synthesis may be contrasted with that schematically illustrated in Figure 1, the prior, conventional method employing DMT protection and separate oxidation and deprotection steps. A further advantage of the invention is illustrated in Figure 3. As shown therein, in Figure 3A, protection and deprotection of hydroxyl groups using DMT is a reversible process, with the DMT cation shown being a relatively stable species. Thus, using DMT as a protecting group can lead to poor yields and unwanted side reactions, insofar as the deprotection reaction is essentially reversible. Figure 3B illustrates the irreversible deprotection reaction of the present invention, wherein nucleophilic attack of the peroxy anion irreversibly cleaves the carbonate moiety, i.e., the *O-p*-chlorophenylcarbonyl group, giving rise to carbon dioxide and the *p*-chlorophenol anion. The reaction is not "reversible," insofar as there is no equilibrium reaction in which a cleaved protecting group could reattach to the hydroxyl moiety, as is the case with removal of DMT.

[0097] As explained earlier herein, the method of the invention also lends itself to synthesis in the 5'-to-3' direction. In such a case, the initial step of the synthetic process involves attachment of a nucleoside monomer to a solid support


at the 5' position, leaving the 3' position available for covalent binding of a subsequent monomer. In this embodiment, i.e., for 5'-to-3' synthesis, a support-bound nucleoside monomer is provided having the structure (IX)

(IX)




wherein O represents the solid support or a support-bound oligonucleotide chain, R is hydrido or hydroxyl, and B is a purine or pyrimidine base. The protected monomer to be added has the structure of formula (X)

(X)



wherein the carbonate protecting group is present at the 3' position, i.e., R¹ is COOR³ where R³ is as defined previously, and R² represents a phosphorus derivative that enables coupling to a free hydroxyl group, preferably a phosphoramidite having the structure (III)

(III)




wherein X and Y are as defined earlier herein. The coupling reaction in which the nucleoside monomer becomes covalently attached to the 3' hydroxyl moiety of the support bound nucleoside is conducted under reaction conditions identical to those described for the 3'-to-5' synthesis. This step of the synthesis gives rise to the intermediate (XI)

5

10

15

(XI)



[0098] As described with respect to oligonucleotide synthesis in the 3'-to-5' direction, the coupling reaction is followed by treatment of the product (XI) with an alpha effect nucleophile in order to remove the carbonate protecting group at the 3' terminus, thus converting the moiety -OR¹ to -OH, and to oxidize the internucleotide phosphite triester linkage to give the desired phosphotriester linkage.

[0099] The two-step process of coupling and deprotection/oxidation is repeated until the oligonucleotide having the desired sequence and length is obtained. Following synthesis, the oligonucleotide may, if desired, be cleaved from the solid support.


[0100] The synthetic methods of the invention may be conducted on any solid substrate having a surface to which chemical entities may bind. Suitable solid supports are typically polymeric, and may have a variety of forms and compositions and derive from naturally occurring materials, naturally occurring materials that have been synthetically modified, or synthetic materials. Examples of suitable support materials include, but are not limited to, polysaccharides such as agarose (e.g., that available commercially as Sepharose®, from Pharmacia) and dextran (e.g., those available commercially under the tradenames Sephadex® and Sephacyl®, also from Pharmacia), polyacrylamides, polystyrenes, polyvinyl alcohols, copolymers of hydroxyethyl methacrylate and methyl methacrylate, silicas, teflons, glasses, and the like. The initial monomer of the oligonucleotide to be synthesized on the substrate surface is typically bound to a linking moiety which is in turn bound to a surface hydrophilic group, e.g., to a surface hydroxyl moiety present on a silica substrate.

[0101] In a related embodiment, the invention features a method for making an oligonucleotide array made up of array features each presenting a specified oligonucleotide sequence at an address on an array substrate. First, the array substrate is treated to protect the hydroxyl moieties on the derivatized surface from reaction with phosphoramidites or analogous phosphorus groups used in oligonucleotide synthesis. Protection involves conversion of free hydroxyl groups to -OR¹ groups, i.e., to carbonate-protected species. The method then involves (a) applying droplets of an alpha effect nucleophile to effect deprotection of hydroxyl moieties at selected addresses and oxidation of the newly formed internucleotide phosphite triester linkages, followed by (b) flooding the array substrate with a medium containing a selected nucleoside monomer having the structure of either Formula (II) (for 3'-to-5' synthesis) or Formula (X) (for 5'-to-3' synthesis). Step (a), deprotection/oxidation, and step (b), monomer addition, are repeated to sequentially build oligonucleotides having the desired sequences at selected addresses to complete the array features. In a variation on the aforementioned method, the applied droplets may comprise the selected nucleoside monomer, while the alpha effect nucleophile is used to flood the array substrate; that is, steps (a) and (b) are essentially reversed.


[0102] In the array construction method according to the invention, the deprotection reagents are aqueous, allowing for good droplet formation on a wide variety of array substrate surfaces. Moreover, because the selection of features employs aqueous media, small-scale discrete droplet application onto specified array addresses can be carried out by adaptation of techniques for reproducible fine droplet deposition from printing technologies.

[0103] The invention additionally provides protected nucleoside monomers as novel compositions of matter useful, *inter alia*, in the synthesis of oligonucleotides as described herein. The novel monomers have the structural formulae (II) and (X)

(II)



(X)



wherein:


B is a purine or pyrimidine base, as described previously herein;

R is hydrido or hydroxyl;

R¹ is COOR³ wherein R³ is as described previously herein, such that the moiety OR¹ represents a carbonate-protected hydroxyl group; and


R² is a phosphorus derivative phosphorus derivative that enables coupling to a free hydroxyl group, and is preferably a phosphoramidite having the structure (III)

(III)




wherein X and Y are as defined earlier herein.

[0104] Reagent (II), used for 3'-to-5' synthesis, is readily prepared by reaction of the unprotected nucleoside with the haloformate R³O-(CO)-Hal wherein Hal represents halo, typically chloro, and R³ is as defined previously, in the presence of a base effective to catalyze the nucleophilic reaction, e.g., pyridine. This step results in a 5'-carbonate, as follows:




[0105] The intermediate so prepared is then phosphitylated with the phosphoramidite $PX_2(OY)$ wherein X and Y are as defined earlier, resulting in conversion of the 3'-hydroxyl moiety to the desired substituent $-O-PX(PY)$, i.e., $-OR^2$:




30 [0106] A specific example of this synthesis is illustrated schematically in Figure 4, wherein "Arco" represents the aryloxycarbonyl group p-chlorophenylcarbonyl, iPr represents isopropyl, and B is either N⁶-benzoyl-protected deoxyadenine, N⁴-Fmoc-protected deoxycytidine, N²-isobutyryl-protected deoxyguanine or thymine. In the initial step of the reaction, the unprotected base is reacted with 4-chlorophenyl chloroformate in the presence of pyridine to give the carbonate-protected 5'-OH, followed by phosphitylation using (iPr₂N)₂PO(CH₂)₂CN, i.e., β-cyanoethyl-N,N-diisopropylamino phosphoramidite.

35 [0107] Reagent (X), used for 5'-to-3' synthesis, may be prepared by first synthesizing a 5'-protected nucleoside using a conventional 5'-OH protecting group such as DMT. This 5'-protected nucleoside is then reacted with the haloformate $R^3O-(CO)-Hal$, which, as above, is done in the presence of a base effective to catalyze the nucleophilic reaction, e.g., pyridine. The DMT group is then removed with acid, resulting in the 3'-carbonate intermediate



55 [0108] Subsequent reaction with the phosphoramidite results in conversion of the 5'-hydroxyl moiety to the desired substituent $-O-PX(PY)$, i.e., $-OR^2$:



[0109] A specific example of this synthesis is illustrated schematically in Figure 5, wherein, as in Figure 4, "Arco" again represents the aryloxy carbonyl group *p*-chlorophenyl carbonyl, *iPr* represents isopropyl, and B is either N⁶-benzoyl-protected deoxyadenine, N⁴-Fmoc-protected deoxycytidine, N²-isobutyryl-protected deoxyguanine or thymine. In the initial step of the reaction shown in Figure 4, the 5'-O-DMT-protected base is reacted with 4-chlorophenyl chloroformate in the presence of pyridine to give the 3' carbonate, followed by DMT removal using trichloroacetic acid and subsequent phosphorylation using β -cyanoethyl-N,N-diisopropylamino phosphoramidite.

[0110] It is to be understood that while the invention has been described in conjunction with the preferred specific embodiments thereof, that the description above as well as the example which follows are intended to illustrate and not limit the scope of the invention. Other aspects, advantages and modifications within the scope of the invention will be apparent to those skilled in the art to which the invention pertains.

[0111] The practice of the present invention will employ, unless otherwise indicated, conventional techniques of synthetic organic chemistry, biochemistry, molecular biology, and the like, which are within the skill of the art. Such techniques are explained fully in the literature.

[0112] The following examples are put forth so as to provide those of ordinary skill in the art with a complete disclosure and description of how to prepare and use the compounds disclosed and claimed herein. Efforts have been made to ensure accuracy with respect to numbers (e.g., amounts, temperature, etc.) but some errors and deviations should be accounted for. Unless indicated otherwise, parts are parts by weight, temperature is in °C and pressure is at or near atmospheric.

[0113] All patents, patent applications, journal articles and other references mentioned herein are incorporated by reference in their entireties.

EXAMPLE 1

PROTECTION AND DEPROTECTION OF DEOXYTHYMIDINE

(A) GENERAL PROCEDURES:

[0114] Nuclear resonance spectra (¹H, ¹³C and ³¹P NMR) were recorded on a Varian VXR-300 spectrometer. Tetramethylsilane was used as an internal reference for ¹H and ¹³C NMR. An external capillary containing 85% H₃PO₄ was used as a reference for ³¹P NMR. Downfield chemical shifts were recorded as positive values for ³¹P NMR. Thin layer chromatography was performed on HF254 silica gel plates (Merck) in: CH₂Cl₂/MeOH, 9:1 (Solvent A), CH₂Cl₂/MeOH, 8:2 (Solvent B), ethyl acetate/THF/Et₃N (45/45/10, v/v/v) (Solvent C). Pyridine, dichloromethane, and benzene were freshly distilled over CaH₂. Acetonitrile was distilled over P₂O₅ (solid), followed by calcium hydride, and stored over molecular sieves. Hexanes and pentanes were distilled. 5'-O-(4,4'-Dimethoxytrityl)-6-N-((di-N-butylamino)methylene)-2'-deoxyadenosine and 2-N-(di-N-butylamino)methylene-2'-deoxyguanosine were prepared according to published procedures. Protected nucleoside-derived CPG was obtained from Applied Biosystems Inc.

(B) SYNTHESIS OF 5'-O-NUCLEOSIDE CARBONATES:

[0115] The syntheses were conducted generally as follows. Deoxythymidine (2 mmol) was co-evaporated with anhydrous pyridine (2 × 20 ml), then redissolved in dry pyridine (40 ml). The corresponding chloroformate (2.2 mmol) was added and the mixture stirred at room temperature (25° C) for 2 hr. The reaction was quenched with water (1 ml), then concentrated. The residual pyridine was removed by co-evaporation with toluene (40 ml).

[0116] The resulting residue was then dissolved in CHCl₃ (50 ml) and extracted with brine (40 ml). The aqueous

layers were back-extracted with CHCl_3 (30 ml). The organic layers were combined, concentrated, and then loaded onto a silica gel column (100 g). The column was eluted with CH_2Cl_2 using a methanol gradient. The isolated products were evaporated to foams.

[0117] This scheme was used to synthesize a series of alkyl and aryl 5'-O-carbonates of deoxythymidine from the corresponding chloroformates. In all cases, the best yields for the 5'-protected nucleoside were obtained when the reactions were performed at room temperature in pyridine using a slight excess of the chloroformate (1.1 eq). Under these conditions, good regioselectivity was observed with most chloroformates. Table 1 sets forth isolated yields of the 5'-protected nucleosides:

TABLE 1

| ISOLATED YIELDS OF 5'-PROTECTED DEOXYTHYMIDINE WITH VARIOUS ALKYL AND ARYL CHLOROFORMATES AT ROOM TEMPERATURE IN PYRIDINE | |
|---|----------------|
| 5'-Carbonate Protected Thymidine | Isolated Yield |
| $\text{Cl}_3\text{C}(\text{CH}_3)_2\text{COCO}_2\text{-dT}$ (1a) [5'-O-TCBOC-dT] | 87 % |
| [5'-O-Fmoc-dT] (1b) | 90 % |
| $2\text{-(NO}_2\text{)}\text{C}_6\text{H}_4\text{OCO}_2\text{-dT}$ (1c) [5'-O-oNPh-dT] | 35 % |
| $\text{C}_6\text{H}_5\text{N=NC}_6\text{H}_4\text{OCO}_2\text{-dT}$ (1d) [5'-O-PAP-dT] | 50 % |
| $\text{C}_6\text{H}_5\text{OCO}_2\text{-dT}$ (1e) [5'-O-Ph-dT] | 60 % |
| $4\text{-(Cl)}\text{C}_6\text{H}_4\text{OCO}_2\text{-dT}$ (1f) [5'-O-pClPh-dT] | 60 % |

[0118] The results were as follows.

[0119] 5'-O-(2,2,2-Trichloro-1,1-Dimethylcarbonyl)Deoxythymidine (5'-O-TCBOC-dT, **1a**):

Yield 87%. R_F (A) = 0.40, R_F (B) = 0.70. $^1\text{H NMR}$ (CDCl_3 +DMSO- D_6) δ : 7.33 (d, 1, H_6), 6.34 (t, J = 7 Hz, 1, H_1), 4.45-4.08 (m, 4, H_3 , H_4 , $\text{H}_{5,5'}$), 2.32-2.1 (m, 2, $\text{H}_{2,2'}$), 1.94-1.93 (m, 6, $\text{C-(CH}_3)_2$), 1.88 (s, 3, $\text{C}^5\text{-CH}_3$). $^{13}\text{C NMR}$ (CDCl_3 +DMSO- D_6) δ : 163.27 (C-4), 150.93 (O-(CO)-O), 149.68 (C-2), 134.21 (C-6), 109.71 (C-5), 104.37 (C- Cl_3), 88.64 (C-Me $_2$), 83.39 (C-4'), 82.94 (C-1'), 62.96 (C-3'), 66.4 (C-5'), 20.02, 19.95 (C-(CH_3) $_2$), 11.6 ($\text{C}^5\text{-CH}_3$).

[0120] 5'-O-(9-Fluorenylmethylcarbonyl)Deoxythymidine (5'-O-Fmoc-dT, **1b**):

Yield 90%. R_F (A) = 0.41, R_F (B) = 0.74. $^1\text{H NMR}$ (CDCl_3 +DMSO- D_6) δ : 7.72-7.28 (m, 9, Fmoc + H_6), 6.36 (t, J = 7 Hz, 1, H_1), 4.54-4.11 (m, 3, CHCH_2 (Fmoc), H_3 , H_4 , $\text{H}_{5,5'}$), 2.35-2.06 (m, 2, $\text{H}_{2,2'}$), 1.79 (s, 3, $\text{C}^5\text{-CH}_3$). $^{13}\text{C NMR}$ (CDCl_3 +DMSO- D_6) δ : 163.87 (C-4), 154.58 (C-2), 150.28 (O-(CO)-O), 142.76, 142.71, 140.91, 127.04, 126.82, 124.59, 119.75 (Fmoc), 134.89 (C-6), 110.55 (C-5), 84.29 (C-4'), 83.76 (C-1'), 69.47 (C-3'), 66.92 (C-5'), 46.3 (Fmoc), 39.86 (C-2'), 12.13 ($\text{C}^5\text{-CH}_3$).

[0121] 5'-O-(α -Nitrophenylcarbonyl)Deoxythymidine (5'-O-oNPh-dT, **1c**):

Yield 35%. R_F (A) = 0.41, R_F (B) = 0.68. $^1\text{H NMR}$ (CDCl_3) δ : 8.21 (d, 1, H_6), 7.89-7.53 (m, 4, aryl), 6.37 (t, J = 7 Hz, 1, H_1), 4.53-4.17 (m, 4, H_3 , H_4 , $\text{H}_{5,5'}$), 2.33-2.03 (m, 2, $\text{H}_{2,2'}$), 1.79 (s, 3, $\text{C}^5\text{-CH}_3$). $^{13}\text{C NMR}$ (CDCl_3) δ : 164.3 (C-4), 153.04 (O-(CO)-O), 152.21 (C-2), 144.68, 142.1, 136.5, 128.36, 126.65, 125.68 (C_6H_4), 136.33 (C-6), 111.05 (C-5), 85.44 (C-1'), 84.62 (C-4'), 71.54 (C-5'), 69.67 (C-3'), 40.15 (C-2'), 12.4 ($\text{C}^5\text{-CH}_3$).

[0122] 5'-O-(p -Phenyldiazophenylcarbonyl)Deoxythymidine (5'-O-PAP-dT, **1d**):

Yield 50%. R_F (A) = 0.41, R_F (B) = 0.75. $^1\text{H NMR}$ (CDCl_3) δ : 7.94-7.28 (m, 10, H_6 +aryl(PAP)), 6.31 (t, J = 7 Hz, 1, H_1), 4.5-4.12 (m, 4, H_3 , H_4 , $\text{H}_{5,5'}$), 2.33-2.19 (m, 2, $\text{H}_{2,2'}$), 1.86 (s, 3, $\text{C}^5\text{-CH}_3$). $^{13}\text{C NMR}$ (CDCl_3) δ : 164.44 (C-4), 152.33 (O-(CO)-O), 152.1 (C-2), 152.86, 152.16, 150.55, 150.23, 131.05, 128.84, 123.86, 122.54, 121.21 (PAP), 135.56 (C-6), 110.92 (C-5), 84.65 (C-1'), 83.55 (C-4'), 70.13 (C-5'), 67.53 (C-3'), 39.73 (C-2'), 11.93 ($\text{C}^5\text{-CH}_3$).

[0123] 5'-O-(Phenylcarbonyl)Deoxythymidine (5'-O-Ph-dT, **1e**):

Yield 60%. R_F (A) = 0.41, R_F (B) = 0.71. $^1\text{H NMR}$ (CDCl_3) δ : 7.54-7.19 (m, 6, H_6 +aryl), 6.34 (t, J = 7 Hz, 1, H_1), 4.52-4.12 (m, 4, H_3 , H_4 , $\text{H}_{5,5'}$), 2.3-2 (m, 2, $\text{H}_{2,2'}$), 1.78 (s, 3, $\text{C}^5\text{-CH}_3$). $^{13}\text{C NMR}$ ($\text{DMSO}-d_6$ +(CD_3) $_2\text{CO}$) δ : 164.36 (C-4), 152.21 (O-(CO)-O), 151.35 (C-2), 154.2, 130.42, 126.97, 121.99 (C_6H_4), 136.61 (C-6), 111.11 (C-5), 85.44 (C-1'), 84.84 (C-4'), 71.73 (C-5'), 68.83 (C-3'), 40.21 (C-2'), 12.5 ($\text{C}^5\text{-CH}_3$).

[0124] 5'-O-(p -Chlorophenylcarbonyl)Deoxythymidine (5'-O-pClPh-dT, **1f**):

Yield 60%. R_F (A) = 0.42, R_F (B) = 0.73. $^1\text{H NMR}$ (CDCl_3) δ : 7.9 (d, 1, H_6), 7.44-7.16 (m, 5, aryl), 6.34 (t, J = 7 Hz, 1, H_1), 4.6-4.12 (m, 4, H_3 , H_4 , $\text{H}_{5,5'}$), 2.3-2.05 (m, 2, $\text{H}_{2,2'}$), 1.74 (s, 3, $\text{C}^5\text{-CH}_3$). $^{13}\text{C NMR}$ (CDCl_3) δ : 164.4 (C-4), 153.23 (O-(CO)-O), 151.4 (C-2), 149.39, 139.86, 129.73, 122.23 (C_6H_4), 136.6 (C-6), 111.1 (C-5), 85.41 (C-1'), 84.8 (C-4'), 71.52 (C-3'), 67.53 (C-5'), 40.25 (C-2'), 12.49 (C^5CH_3).

(C) SYNTHESIS OF 5'-O-DMT-3'-O-R-DEOXYTHYMIDINES:

[0125] The 3'-hydroxyl group of 5'-O-DMT-deoxythymidine was protected with phenyloxycarbonyl (**2a**), benzoyl (**2b**), and acetyl (**2c**), as follows. 5'-O-(4,4'-Dimethoxytrityl)-deoxythymidine (1 mmol) was co-evaporated 3 times with anhydrous pyridine, then redissolved in 20 ml of pyridine. Corresponding chloroformates (1.1 mmol) were added to the nucleoside mixture. After stirring for 6 hr, the reaction was quenched with water (100 ml) and concentrated. Residues of pyridine were removed by co-evaporation with toluene (2 × 20 ml).

[0126] The resulting gum was dissolved in CH₂Cl₂, extracted with 10% aqueous NaHCO₃, and dried over Na₂SO₄. After concentration, the product was loaded onto a silica gel column (50 g) and eluted with CH₂Cl₂ using a methanol gradient (0-3%). Product fractions were collected and concentrated to a foam.

[0127] The results were as follows.

[0128] 5'-O-(4,4'-Dimethoxytrityl)-3'-O-Phenylcarbonyl Deoxythymidine (**2a**):

Yield 80%. R_F (A) = 0.74, R_F (B) = 0.91. ¹H NMR (CDCl₃) δ: 7.65-6.83 (m, 18, H₆+DMTr+aryl), 6.57 (t, J = 7 Hz, 1, H₁), 5.45 (m, 1, H₃), 4.34 (m, 1, H₄), 3.79 (s, 6, OCH₃), 3.54 (m, 2, H_{5,5'}), 2.72-2.52 (m, 2, H_{2,2'}), 1.41 (s, 3, C⁵-CH₃).

[0129] 5'-O-(4,4'-Dimethoxytrityl)-3'-O-Benzoyl Deoxythymidine (**2b**):

Yield 90%. R_F (A) = 0.72, R_F (B) = 0.91. ¹H NMR (CDCl₃) δ: 8.07-6.85 (m, 18, H₆+DMTr+aryl), 6.58 (t, J = 7 Hz, 1, H₁), 5.45 (m, 1, H₃), 4.14 (m, 1, H₄), 3.79 (s, 6, OCH₃), 3.57 (m, 2, H_{5,5'}), 2.63 (m, 2, H_{2,2'}), 1.42 (s, 3, C⁵-CH₃).

[0130] 5'-O-(4,4'-Dimethoxytrityl)-3'-O-Acetyl Deoxythymidine (**2c**):

Yield 90%. R_F (A) = 0.67, R_F (B) = 0.89. ¹H NMR (CDCl₃) δ: 7.62 (s, 1, H₆), 7.4-6.82 (m, 13, DMTr), 6.46 (t, J = 7 Hz, 1, H₁), 5.45 (m, 1, H₃), 4.14 (m, 1, H₄), 3.78 (s, 6, OCH₃), 3.47 (m, 2, H_{5,5'}), 2.45 (m, 2, H_{2,2'}), 2.08 (s, 3, CO-CH₃), 1.39 (s, 3, C⁵-CH₃).

(D) NUCLEOSIDE DEPROTECTION BY PEROXY ANIONS:

[0131] Deprotection reactions were carried out using peroxy anions on alkyl and aryl 5'-O-carbonates of deoxythymidine synthesized as described above. The reactions were monitored by TLC for complete conversion of the starting material to deoxythymidine. A wide variety of peroxy anions, known to exhibit strong alpha effects, were screened for their ability to cleave 5'-O-carbonates of deoxythymidine. Peroxy anion solutions active in cleavage of the 5'-O-carbonates were buffered at a variety of pH conditions. The cleavage activity of these peroxy anion solutions was shown to be rapid only at pH conditions above the pK_a for the formation of the anion. The ability of peroxy anion solutions **A**, **B**, **C**, **D** and **E** to completely deprotect the 5'-O-carbonates of deoxythymidines **1a** - **1f** is summarized in Table 2.

Solution A: 3.1% LiOH•H₂O (10 mL), 1.5 M 2-amino-2-methyl-1-propanol ("AMP"), pH 10.3 (15 mL), 1,4-dioxane (50 mL), 30% H₂O₂ (12 mL), pH 12.0.

Solution B: 3.1% LiOH•H₂O (10 mL), 1.5 M 2-amino-2-methyl-1-propanol ("AMP"), pH 10.3 (15 mL), dimethyl sulfoxide ("DMSO") (50 mL), 30% H₂O₂ (12 mL), pH 12.0.

Solution C: 3.1% LiOH•H₂O (10 mL), 1.5 M 2-amino-2-methyl-1-propanol ("AMP"), pH 10.3 (15 mL), 1,4-dioxane (50 mL), 30% H₂O₂ (12 mL), pH 12.0, m-chloroperbenzoic acid ("mCPBA") (1.78 g), pH 9.6.

Solution D: H₂O (10 mL), dioxane (50 mL), 2.5 M Tris (15 mL), H₂O₂ (12 mL), mCPBA (1.78 g), pH 9.0.

Solution E: H₂O (10 mL), dioxane (50 mL), 2.5 M Tris (15 mL), t-butyl-OOH (0.1 M), pH 9.0.

TABLE 2

| TIMES REQUIRED FOR COMPLETE CONVERSION OF PROTECTED NUCLEOSIDES 1A THROUGH 1F USING PEROXY ANION SOLUTIONS A, B, C, D AND E | | | | | |
|---|--|---------|----------|---------|---------|
| 5'-Carbonate-dT Compounds | Reaction Completion Times for Deprotection Solutions | | | | |
| | A | B | C | D | E |
| 1a | < 1 min | < 1 min | < 12 min | - | - |
| 1b | > 1 hr | < 1 min | > 3 hr | - | - |
| 1c | < 1 min | < 1 min | < 1 min | - | - |
| 1d | < 1 min | < 1 min | < 1 min | < 1 min | > 12 hr |
| 1e | < 1 min | < 1 min | < 1 min | < 2 min | < 12 hr |

TABLE 2 (continued)

| TIMES REQUIRED FOR COMPLETE CONVERSION OF PROTECTED NUCLEOSIDES 1A THROUGH 1F USING PEROXY ANION SOLUTIONS A, B, C, D AND E | | | | | |
|---|--|---------|---------|---------|---------|
| | Reaction Completion Times for Deprotection Solutions | | | | |
| 5'-Carbonate-dT Compounds | A | B | C | D | E |
| 1f | < 1 min | < 1 min | < 1 min | < 1 min | < 12 hr |

(E) SELECTIVITY OF VARIOUS PEROXY ANION SOLUTIONS FOR DEPROTECTION OF CARBONATES:

[0132] As described in part (C) of this example, the 3'-hydroxyl group of 5'-O-DMT-deoxythymidine was protected with a phenyloxycarbonyl (**2a**), a benzoyl (**2b**), and an acetyl (**2c**) group. The stability of these 3'-protecting groups was determined by TLC using deprotection conditions **C** and **D** (Table 2). Under both these conditions, the phenyl carbonate was completely removed in less than 2 min. The 3'-benzoyl group was completely stable under both conditions for 140 min. The 3'-acetyl group was cleaved to a small extent (less than 3%) over the 140 min exposure to deprotection condition **A** (pH 10.0). The 3'-benzoyl group was completely stable for the 140 min exposure to condition **B**.

(F) SELECTIVITY OF DEPROTECTION ON SOLID-SUPPORT ATTACHED NUCLEOSIDES:

[0133] The demonstration of stability at the 3' position was then extended to the succinate linker commonly used for the attachment of nucleosides to Controlled Pore Glass, as follows. 5'-DMT-deoxythymidine attached to Long Chain Alkyl Amine Controlled Pore Glass (LCAA/CPG) through a 3'-succinate linkage was obtained from a commercial source. This solid-support attached nucleoside was then exposed to deprotection conditions **A** through **D**. The stability of the 3'-linkage was determined spectrophotometrically based upon the evolution of the trityl cation during subsequent treatment with toluene sulfonic acid in anhydrous acetonitrile. Deprotection conditions **A** and **B** gave complete cleavage of the 3'-succinate in 20 min. Deprotection conditions **C** and **D** gave less than 2% cleavage of the 3'-succinate after 20 hrs.

EXAMPLE 2**SIMULTANEOUS OLIGOTHYMIDYLATE DEPROTECTION AND INTERNUCLEOTIDE BOND OXIDATION BY PEROXY ANIONS****(A) OLIGONUCLEOTIDE SYNTHESIS ON CONTROLLED PORE GLASS:**

[0134] Oligonucleotides were synthesized on CPG using an automated DNA synthesizer (ABI model 380A). The synthesis cycle used for 5'-DMT protected nucleoside phosphoramidites (Cycle 1) is shown in Table 4. This cycle was initially modified for the use of 5'-carbonate protected nucleoside phosphoramidites simply by substituting the alternative deprotection mixtures for the 3% TCA solution (Step 8, Table 4) and varying the exposure times. For the synthesis of longer sequences using 5'-carbonate protected nucleoside phosphoramidites, it was necessary to separate the deprotection mixture into a two-component system (Table 3). The separation of the deprotection mixture was accomplished using the capping ports on the synthesizer, and thus necessitated elimination of the capping step from the synthesis cycle. Table 4 shows the optimized cycle for synthesis using 5'-carbonate protected nucleoside phosphoramidites (Cycle 2):

TABLE 3

| TWO-COMPONENT SYSTEM FOR STORAGE OF DEPROTECTION SOLUTION C | |
|---|---|
| Solution C-1 | 30 % H ₂ O ₂ (10 ml), LiOH (280 mg), dioxane (7.5 ml), 2.5 M Tris-Base (15 ml), water (42.5 ml) |
| Solution C-2 | 50 - 60 % mCPBA (1.78 g), dioxane (42.5 ml) |

TABLE 4

| OLIGONUCLEOTIDE SYNTHESIS CYCLES | | | | |
|----------------------------------|-----------|--|--------------------|--------------------|
| Step # | Function | Reagent | Cycle 1 Time, sec. | Cycle 2 Time, sec. |
| 1 | Wash | Acetonitrile | 25 | 25 |
| 2 | Coupling | Amidite (0.15 M, 30 eq) Tetrazole (0.5 M, 120 eq) in Anhydrous Acetonitrile | 2 × 30 | 2 × 30 |
| 3 | Wash | Acetonitrile | 5 | 5 |
| 4 | Capping | N-Methylimidazole/2,6-Lutidine/Acetic Anhydride/THF (1/1/1/2, vol/vol/vol/vol) | 40 | - |
| 5 | Oxidation | 0.1 M I ₂ in THF/Lutidine/Water (80/40/2, vol/vol/vol) | 30 | - |
| 6 | Wash | Acetonitrile | 25 | - |
| 7 | Wash | Dichloromethane (Cycle 1) 1,4-Dioxane (Cycle 2) | 25 | 25 |
| 8 | Deblock | 3 % TCA in CH ₂ Cl ₂ (Cycle 1) 1:1 mix of Solution C-1 & Solution C-2 from Table 3 (Cycle 2) | 2 × 30 | 480 |
| 9 | Wash | Dichloromethane (Cycle 1) 1,4-Dioxane (Cycle 2) | 25 | 25 |

(B) ANALYSIS OF OLIGONUCLEOTIDES BY HPLC:

[0135] The oligonucleotides synthesized on the solid support were deprotected with concentrated ammonium hydroxide (55° C, 24 hr). The ammonium hydroxide solutions were removed from the support and evaporated to dryness. The crude oligonucleotides were reconstituted in distilled water and stored at -20° C.

[0136] HPLC analysis was performed by ion-exchange HPLC (Nucleogen 60-7DEAE, 4 mm ID x 125 mm). Oligonucleotides were eluted from the column with a LiCl gradient (0.0-0.7 M) in a water/acetonitrile (60/40, v/v) buffer containing sodium acetate (0.002 M, pH 6.0).

(C) SOLID-SUPPORT DEPROTECTION OF 5'-O-CARBONATES OF THYMIDINE:

[0137] The deprotection efficiency of peroxy-anion solutions on oligonucleotides was determined by the synthesis of oligothymidylate tetramers. The 5'-O-arylcarbonates of deoxythymidine (see part (B) of Example 1, compounds **1a** through **1f**) were converted to the corresponding 3'-O-(2-cyanoethyl)-N,N-diisopropylphosphoramidite by procedures described generally in A.D. Barone *et al.* (1984) *Nucleic Acids Res.* **12**:4051, as follows.

[0138] Synthesis of the 2-cyanoethyl-N,N,N',N'-tetraisopropyl-phosphorodiamidite phosphine was performed according the procedure described in A. Kraszewski *et al.* (1987) *Nucleic Acids Res.* **15**:177. The resulting product was purified by distillation from CsF. The product was obtained in 60% yield. Purity was confirmed by ³¹P NMR (CDCl₃) δ: 123.8 ppm.

[0139] Thymidyl-3'-5'-deoxythymidine was synthesized on solid-support using 5'-O-dimethoxytrityl-3'-O-(2-cyanoethyl)-N,N-diisopropylaminodeoxythymidinephosphoramidite. The dimer was elongated to a trimer using a 5'-O-aryloxycarbonyl-3'-O-(2-cyanoethyl)-N,N-diisopropylaminodeoxythymidinephosphoramidite and synthesis cycle 1 (Table 4). Deprotection of the carbonate was then attempted using deprotection mixture C, at 1 min increments, from 1-15 min. The extent of deprotection was determined by the yield of the subsequent coupling reaction using a standard 5'-DMT-dT phosphoramidite. Deprotection efficiency for the 5'-O-arylcarbonate was determined using ion-exchange HPLC. The percent deprotection was calculated by integration and normalization of peak areas for the corresponding trimers and tetramers, assuming quantitative coupling reactions. The optimum deprotection time and extent of deprotection for each aryloxycarbonyl group is summarized in Table 5.

TABLE 5

| OPTIMUM DEPROTECTION TIMES DETERMINED FOR 5'-ARYLCARBONATES OF THYMIDINE ON CONTROLLED PORE GLASS USING DEPROTECTION SOLUTION C | | |
|---|---------------------------|-------------------------|
| 5'-Carbonate dT Compounds | Optimum Deprotection Time | Deprotection Efficiency |
| 1c | 5 min | 80 % |
| 1d | 1 min | 94 % |
| 1e | 7 min | 98 % |
| 1f | 3 min] | 98 % |

(D) SOLID SUPPORT SYNTHESIS AND INTERNUCLEOTIDE BOND OXIDATION:

[0140] Several oligothymidylate tetramers were synthesized on Controlled Pore Glass using 5'-O-*p*-chlorophenylloxycarbonyl-3'-O-(2-cyanoethyl)-N,N-diisopropylaminodeoxythymidine-phosphoramidite. These syntheses were performed on a 1 μ mol scale using an automated DNA synthesizer. The only modification from the standard 1 μ mol synthesis cycle (Cycle 1, Table 4) was the use of deprotection mixture **C** (7 min) in place of 3% TCA in dichloromethane. The resulting tetramers were compared to oligothymidylate tetramers synthesized using the standard DMT protected phosphoramidites of thymidine. These tetramers were then analyzed using ion-exchange HPLC. There were no detectable differences in the yield or purity of any of the oligomers.

[0141] Oligothymidylate tetramers were then synthesized using this same synthesis cycle, which was again modified by the removal of the iodine oxidation step. This concomitant deprotection and oxidation cycle produced tetramers of identical yield and purity to the standard DMT phosphoramidite synthesis. Decomposition of MCPBA in the presence of LiOH results in the deprotection mixture being effective for only a few hours. In order to synthesize longer sequences, it was necessary to separate the deprotection mixture into a two component system (Table 3). This was accomplished using the capping ports on the automated DNA synthesizer. Separating the LiOH from the mCPBA and mixing just prior to deprotection allows the reagents to remain effective for several days. Oligonucleotide synthesis using 5'-O-arylcarbonate nucleoside phosphoramidites was carried out with and without acetic anhydride capping. No adverse effects on the yield of final product or increases in the appearance of n-1 products were observed in absence of capping. This is contrary to what is seen with the use of DMT protected phosphoramidites in the absence of capping. Anion-exchange HPLC profiles of crude synthesis products of oligothymidylate decamers were produced. Product purity and yield of full-length oligonucleotides, using peroxyanion deprotection of 5'-O-carbonates in absence of acetic anhydride capping and iodine oxidation (Cycle 2, Table 4), were comparable to or better than those obtained using DMT phosphoramidites and the standard synthesis cycle.

EXAMPLE 3**PEROXY ANION DEPROTECTION OF 5'-O-DMT-PROTECTED CYTOSINE, ADENINE, URACIL, THYMIDINE AND GUANOSINE NUCLEOSIDES**

[0142] The unprotected heterocyclic bases cytosine and adenine are susceptible to N-oxidation by peracids and peroxides under stringent conditions, and oxidative reactions that result in ring cleavage of uracil, thymidine and guanosine in the presence of highly concentrated peroxides at elevated temperatures have been described. 5'-O-DMT-protected nucleosides, N-protected with a (di-N-butylamino)methylene group, were dissolved in deprotection mixture **C** and allowed to react for 24 hrs. The tritylated nucleosides were extracted from the aqueous deprotection mixture with CHCl_3 and analyzed by ^{13}C NMR and TLC. Neither formation of N-oxides nor attack at the 5,6-double bond of thymidine (leading to ring cleavage) was detected.

EXAMPLE 4**SYNTHESIS OF MIXED OLIGONUCLEOTIDES**

[0143] This example demonstrates extension of the method of the invention to synthesis of mixed oligonucleotide sequences, employing substituted aryl carbonate protected phosphoramidite synthons, and following each coupling reaction by treatment with a mixture of peroxy-anions at mild pH (less than 10) to deprotect and concomitantly oxidize the internucleotide linkage.

[0144] The method is high-yielding, and effective for the four main 2'-deoxynucleotides. Synthesis in both the 3' - 5' direction and the 5' - 3' direction were carried out, with equal effect.

(A) PROTECTED PHOSPHORAMIDITE SYNTHESIS:

[0145] Generally, the protected nucleoside phosphoramidites were prepared as follows. The 3'- or 5'-protected nucleoside (5.00 mmol) and tetrazole (175 mg, 2.50 mmol) were dried under vacuum for 24h and then dissolved in trichloromethane (100 mL). 2-cyanoethyl-*N,N,N',N'*-tetraisopropyl phosphane (2.06 mL, 6.50 mmol) was added in one portion and the mixture stirred over 1 hour. The reaction mixture was washed with sat. NaHCO_3 (150 mL) and brine (150 mL), dried over MgSO_4 and applied directly to the top of a silica column equilibrated with hexanes. The dichloromethane was flashed off the column with hexanes, and the product eluted as a mixture of diastereoisomers using 1/1 hexanes/ethyl acetate then ethyl acetate. After evaporation of solvents *in vacuo* and coevaporation with dichloromethane, products were isolated as friable, white, glassy solids in yields varying from 70% to 90%.

[0146] The four 5'-aryloxycarbonyl-3'-nucleoside phosphoramidites were prepared by the straightforward two-step procedure shown generally in Figure 4. In a first step, commercially available base protected 2'-oligodeoxynucleosides were selectively aryl carbonate protected at the 5' position by treatment with 4-chlorophenyl chloroformate in dilute anhydrous pyridine to yield 5'-aryloxycarbonyl protected compounds in moderate to good yield. The use of more concentrated reaction mixtures resulted in an increase in the amounts of isolated 3'- and 3', 5'-*bis*-aryloxycarbonyl-protected materials. In a second step, the resulting compounds were phosphitylated using the method described in Barone *et al., supra*, to furnish high yields following column chromatography.

[0147] Synthesis of the four 3'-aryloxycarbonyl-5'-nucleoside phosphoramidites were prepared by the three-step procedure shown in Figure 5.

(C) DEPROTECTION MIXTURE:

[0148] The deprotection mixture was formulated in two parts, which were mixed immediately prior to use. Solution F: 3.1% w/v lithium hydroxide monohydrate (10 mL), 1.5 M 2-amino-2-methyl-1-propanol pH 10.3 (15 mL), 1,4 dioxane (17.5 mL). Solution G: 1,4-dioxane (32.5 mL), 50 - 83% 3-chloroperbenzoic acid (1.78 g), 30% hydrogen peroxide (12 mL). The initial pH of the deprotection mixture was 9.6 ± 0.05 . For pH dependence studies, the initial deprotection mixture was altered by varying the strength of the lithium hydroxide solution.

(D) SYNTHESIS OF MIXED-SEQUENCE OLIGONUCLEOTIDES:

[0149] A series of model oligodeoxynucleotides was synthesized, having sequences 3'-T₃AT₂AT₃-5', 3'-T₃CT₂CT₃-5', 3'-T₃GT₂GT₃-5', 3'-TACGT-5', 3'TACGTACGT-5', 3'-TA₇T-5', 5'-TACGT-3', 5'-TACGTACGT-3', and 5'-CAGTTGTAAACGAGTT-3'. HPLC analysis was performed as described in Example 2, part (B); HPLC traces of the all products confirmed the results.

[0150] The HPLC obtained for 5'-CAGTTGTAAACGAGTT-3' is shown in Figure 6. The calculated molecular weight for 5'-CAGTTGTAAACGAGTT-3' is 4921.1; the actual molecular weight determined using MALDI (Matrix Absorption Laser Desorption Ionization) TOF (Time of Flight) analysis was 4921.9. The MALDI TOF spectrum is shown in Figure 7.

(E) STABILITY OF BASE PROTECTING GROUPS IN THE DEPROTECTION MIXTURE:


[0151] The stability of the standard base protecting groups A^{Bz}, C^{Bz}, and G^{ibu} during exposure to the deprotection mixture was tested by incubating 5'-DMT base-protected deoxynucleosides at room temperature with a large excess of the deprotection mixture. The extent of cleavage of the base protecting groups over time was measured by TLC. The approximate T_{1/2} values for A^{Bz}, C^{Bz}, and G^{ibu} were approximately ½ hour, 2 hours, and 1 day, respectively, and unlikely to present difficulties for syntheses.

Claims

1. A method for synthesizing oligonucleotides which comprises (a) condensing the 3'-OH or 5'-OH group of a support-bound nucleoside or oligonucleotide with a monomeric nucleoside phosphoramidite having a carbonate-protected hydroxyl group, to provide an intermediate in which the support-bound nucleoside or oligonucleotide is bound to the monomeric nucleoside through a phosphite triester linkage, and (b) treating the intermediate provided in step (a) with a deprotecting reagent effective to convert the carbonate-protected hydroxyl group to a free hydroxyl moiety and simultaneously oxidize the phosphite triester linkage to give a phosphotriester linkage

2. The method of claim 1, wherein step (b) is conducted at neutral to mildly basic pH.
3. The method of claim 1 or 2 wherein the nucleophile is a peroxide.
- 5 4. The method of any preceding claim, wherein synthesis is conducted in the 3'-to-5' direction, and step (a) involves condensation at the 5'-OH of the support-bound nucleoside or oligonucleotide.
- 10 5. The method of any preceding claim, wherein the monomeric nucleoside phosphoramidite has the structure of formula (II)

(II)



wherein:


B is a purine or pyrimidine base;

R is hydrido or hydroxyl;

R¹ is COOR³, wherein R³ is substituted or unsubstituted hydrocarbyl;

R² has the structure (III)


(III)




wherein X is NQ¹Q² in which Q¹ and Q² are independently selected from the group consisting of alkyl, aryl, aralkyl, alkaryl, cycloalkyl, alkenyl, cycloalkenyl, alkynyl, or cycloalkynyl, optionally containing one or more nonhydrocarbyl linkages and optionally substituted on one or more available carbon atoms with a nonhydrocarbyl substituent, or wherein Q¹ and Q² are linked to form a mono- or polyheterocyclic ring, and Y is hydrido or hydrocarbyl.

6. The method of claim 5, wherein R³ comprises a moiety that becomes fluorescent or coloured upon cleavage of the carbonate -OCOOR³.
7. A method for making an oligonucleotide array made up of array features each presenting a specified oligonucleotide sequence at an address on an array substrate, the method comprising steps of:
 - providing a hydroxyl-derivatized array substrate and treating the array substrate to protect hydroxyl moieties on the derivatized surface from reaction with phosphoramidites,
 - then iteratively carrying out the steps of (i) applying droplets of an alpha effect nucleophile to effect deprotection of hydroxyl moieties at selected addresses, and (ii) flooding the array substrate with a medium containing a selected monomeric nucleoside phosphoramidite having a carbonate-protected hydroxyl group, to permit covalent attachment of the selected nucleoside to the deprotected hydroxyl moieties at the selected addresses.
8. A nucleoside monomer having the structure of formula (II) or formula (X)

(II)




(X)



wherein:

B is a purine or pyrimidine base, as described previously herein;
 R is hydrido or hydroxyl;
 R¹ is COOR³, wherein R³ is substituted or unsubstituted hydrocarbyl; and
 R² has the structure (III)

(III)



wherein X is NQ¹Q² in which Q¹ and Q² are independently selected from the group consisting of alkyl, aryl, aralkyl, alkaryl, cycloalkyl, alkenyl, cycloalkenyl, alkynyl, or cycloalkynyl, optionally containing one or more nonhydrocarbonyl linkages and optionally substituted on one or more available carbon atoms with a nonhydrocarbonyl substituent, or wherein Q¹ and Q² are linked to form a mono- or polyheterocyclic ring, and Y is hydrido or hydrocarbonyl.

9. A nucleoside monomer as claimed in claim 8 wherein R³ comprises a moiety that becomes fluorescent or coloured upon cleavage of the carbonate -OCOOR³.
10. A kit for synthesizing an oligonucleotide on a solid support, comprising a hydroxyl-derivatized support surface, a protecting group for protecting hydroxyl moieties on said derivatized support surface, at least one protected nucleoside, at least one nucleoside phosphoramidite, a nucleophile that exhibits an alpha effect at neutral to mildly basic pH, and reagents suitable for establishing pH and for carrying out reactions of deprotection, phosphoramidite coupling, and oxidation to form an internucleotide phosphotriester linkage.

Figure 1

3'-5' Synthesis With DMT Protection





Figure 2

3'-5' Synthesis With Carbonate Protection



○ = Solid support/growing oligonucleotide




Figure 3
Comparison of Deprotection Mechanisms

Figure 3A:




Figure 3B:





Figure 4

5'-Arco-3'-Amidite Synthesis




B =

a A(N⁶-Bz)b C(N⁴-Fmoc)c G(N²-ibu)

d T

Arco =




(i) 4-Chlorophenyl chloroformate/pyridine/2 h;

(ii) (iPr₂N)₂PO(CH₂)₂CN/tetrazole/CH₂Cl₂/1 h.

Figure 5

5'-Arco-3'-Amidite Synthesis



B =

a A(N⁶-Bz)b C(N⁴-Fmoc)c G(N²-ibu)

d T

(i) 4-Chlorophenyl chloroformate/pyridine/2 h;

(ii) 3% Trichloroacetic acid/CH₂Cl₂/3 min;(iii) ((iPr₂N)₂PO(CH₂)₂CN/tetrazole/CH₂Cl₂/1 h.

Figure 6





Figure 7



BEST AVAILABLE COPY



(11) **EP 0 984 021 A3**

(12) **EUROPEAN PATENT APPLICATION**

(88) Date of publication A3:
01.08.2001 Bulletin 2001/31

(51) Int Cl.7: **C07H 21/00, C07B 61/00,
C07H 19/10, C07H 19/20**

(43) Date of publication A2:
08.03.2000 Bulletin 2000/10

(21) Application number: **99306168.8**

(22) Date of filing: **03.08.1999**

(84) Designated Contracting States:
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE**
Designated Extension States:
AL LT LV MK RO SI

(30) Priority: **03.08.1998 US 128052
22.06.1999 US 338179**

(71) Applicant: **Agilent Technologies Inc.
Palo Alto, CA 94303 (US)**

(72) Inventors:
• **Dellinger, Douglas J.
Sunnyvale, CA 94087 (US)**
• **Caruthers, Marvin H.
Boulder, CO 80303 (US)**
• **Betley, Jason R.
Bury St. Edmunds, Suffolk IP28 6HW (GB)**


(74) Representative: **Tollett, Ian et al
Williams, Powell & Associates,
4 St. Paul's Churchyard Inn
London EC4M 8AY (GB)**

(54) **Synthesis of oligonucleotides**

(57) The invention provides a method for synthesizing oligonucleotides using carbonate protection of hydroxyl groups and nucleophilic deprotection reagents. The deprotection reagents irreversibly cleave the carbonate protecting groups while simultaneously oxidizing the internucleotide phosphite triester linkage, and can be used in aqueous solution at neutral to mildly basic pH. The method eliminates the need for separate deprotection and oxidation steps, and, since the use of acid to remove protecting groups is unnecessary, acid-induced depurination is avoided. Fluorescent or other readily detectable carbonate protecting groups can be used, enabling monitoring of individual reaction steps during oligonucleotide synthesis. The invention is particularly useful in the highly parallel, microscale synthesis of oligonucleotides. Reagents and kits for carrying out the aforementioned method are provided as well.

Figure 1

3'-5' Synthesis With DMT Protection



EP 0 984 021 A3



European Patent
Office

PARTIAL EUROPEAN SEARCH REPORT

Application Number

which under Rule 45 of the European Patent Convention shall be considered, for the purposes of subsequent proceedings, as the European search report

EP 99 30 6168

| DOCUMENTS CONSIDERED TO BE RELEVANT | | | |
|--|---|---|--|
| Category | Citation of document with indication, where appropriate, of relevant passages | Relevant to claim | CLASSIFICATION OF THE APPLICATION (Int.C1.7) |
| X | WO 96 28457 A (DUKE UNIVERSITY) 19 September 1996 (1996-09-19) * page 31, compound 4 * | 8,9 | C07H21/00 C07B61/00 C07H19/10 C07H19/20 |
| X | EP 0 742 287 A (MCGALL ET AL.) 13 November 1996 (1996-11-13) * page 16 - page 17, examples 9-11 * | 8,9 | |
| X | HASAN ET AL.: "BASE-BORONATED DINUCLEOTIDES: SYNTHESIS AND EFFECT OF N7-CYANOBORANE SUBSTITUTION ON THE BASE PROTONS" NUCLEIC ACID RESEARCH, vol. 24, no. 11, 1996, pages 2150-2157, XP002167837 * compounds 6-9, page 2151, column 2 - page 2152, column 1 * | 8,9 | |
| | | | TECHNICAL FIELDS SEARCHED (Int.C1.7) |
| | | | C07H C07B |
| INCOMPLETE SEARCH | | | |
| <p>The Search Division considers that the present application, or one or more of its claims, does/does not comply with the EPC to such an extent that a meaningful search into the state of the art cannot be carried out, or can only be carried out partially, for these claims.</p> <p>Claims searched completely :</p> <p>Claims searched incompletely :</p> <p>Claims not searched :</p> <p>Reason for the limitation of the search:</p> <p>see sheet C</p> | | | |
| Place of search | | Date of completion of the search | Examiner |
| THE HAGUE | | 28 May 2001 | Fitz, W |
| CATEGORY OF CITED DOCUMENTS | | <p>T : theory or principle underlying the invention</p> <p>E : earlier patent document, but published on, or after the filing date</p> <p>D : document cited in the application</p> <p>L : document cited for other reasons</p> <p>& : member of the same patent family, corresponding document</p> | |
| <p>X : particularly relevant if taken alone</p> <p>Y : particularly relevant if combined with another document of the same category</p> <p>A : technological background</p> <p>O : non-written disclosure</p> <p>P : intermediate document</p> | | | |

EPO FORM 1503 03 89 (P04C07)



European Patent
Office

INCOMPLETE SEARCH
SHEET C

Application Number
EP 99 30 6168

Claim(s) searched completely:

3

Claim(s) searched incompletely:

1,2,4-10

Reason for the limitation of the search:

Present claims 1,2,4-7,10 relate to a method containing a reagent defined by reference to a desirable characteristic or property, namely a deprotecting reagent effective to convert the carbonate-protected hydroxyl group to a free hydroxyl moiety and simultaneously oxidize the phosphite triester linkage to give a phosphotriester linkage.

The claims cover all methods containing this reagent having this characteristic or property, whereas the application provides support within the meaning of Article 84 EPC and/or disclosure within the meaning of Article 83 EPC for only a very limited number of such reagents. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Independent of the above reasoning, the claims also lack clarity (Article 84 EPC). An attempt is made to define the reagent by reference to a result to be achieved. Again, this lack of clarity in the present case is such as to render a meaningful search over the whole of the claimed scope impossible. Consequently, the search has been carried out for those parts of the claims which appear to be clear, supported and disclosed. Thus the above claims have only been completely searched in the light of claim 3 i.e. where the reagent is a peroxide.

Furthermore with regard to claims 8 and 9, the initial phase of the search revealed a very large number of documents relevant to the issue of novelty. So many documents were retrieved that it is impossible to determine which parts of the claims may be said to define subject-matter for which protection might legitimately be sought (Article 84 EPC). For these reasons, a meaningful search over the whole breadth of the claim(s) is impossible. Consequently, the search has been restricted to:

compounds given in the Formula of claim 8, where R3 = aryloxycarbonyl.



European Patent
Office

PARTIAL EUROPEAN SEARCH REPORT

Application Number
EP 99 30 6168

| DOCUMENTS CONSIDERED TO BE RELEVANT | | | CLASSIFICATION OF THE APPLICATION (Int.Cl.7) |
|-------------------------------------|---|-------------------|--|
| Category | Citation of document with indication, where appropriate, of relevant passages | Relevant to claim | |
| X | BERGMANN ET AL.: "ALLYL AS INTERNUCLEOTIDE PROTECTING GROUP IN DNA SYNTHESIS TO BE CLEAVED OFF BY AMMONIA" TETRAHEDRON, vol. 51, no. 25, 1995, pages 6971-6976, XP002167838 * compound 1e, page 6972 * | 8,9 | |
| X | BERGMANN F ET AL: "NUCLEOTIDES PART XLI 1) THE 2-DANSYLETHOXYCARBONYL (=2-((5-DIMETHYLAMINO)NAPHTHALEN-1-YL)SULFONYL)ETHOXYCARBONYL;DNSEOC) GROUP FOR PROTECTION OF THE 5'-HYDROXY FUNCTION IN OLIGORIBONUCLEOTIDE SYNTHESIS" HELVETICA CHIMICA ACTA, vol. 77, 1994, pages 203-214, XP002167878 ISSN: 0018-019X * compounds 26-33, page 205, scheme 2; page 206, lines 20-21 * | 8,9 | TECHNICAL FIELDS SEARCHED (Int.Cl.7) |
| X | PIRRUNG ET AL.: "PROOFING OF PHOTOLITHOGRAPHIC DNA SYNTHESIS WITH 3',5'-DIMETHOXYBENZOINYLOXYCARBONYL-PROTECTED DEOXYNUCLEOSIDE PHOSPHORAMIDITES" JOURNAL OF ORGANIC CHEMISTRY, vol. 63, 1998, pages 241-246, XP000916248 * compounds 1-4, page 242, chart 1; page 243, figure 2 * | 8,9 | |
| X | SIGMUND ET AL.: "A NEW TYPE OF FLUORESCENCE LABELLING OF NUCLEOSIDES, NUCLEOTIDES AND OLIGONUCLEOTIDES" NUCLEOSIDES AND NUCLEOTIDES, vol. 16, 1997, pages 685-696, XP002167901 * page 687, compound 13 * | 8,9 | |
| | -/-- | | |

EPO FORM 1503 01.82 (P04C10)



European Patent
Office

PARTIAL EUROPEAN SEARCH REPORT

Application Number
EP 99 30 6168

| DOCUMENTS CONSIDERED TO BE RELEVANT | | | CLASSIFICATION OF THE APPLICATION (Int.Cl.7) |
|-------------------------------------|--|-------------------|--|
| Category | Citation of document with indication, where appropriate, of relevant passages | Relevant to claim | |
| X | MCGALL G H ET AL.: "THE EFFICIENCY OF LIGHT-DIRECTED SYNTHESIS OF DNA ARRAYS ON GLASS SUBSTRATES" JOURNAL OF THE AMERICAN CHEMICAL SOCIETY, US, AMERICAN CHEMICAL SOCIETY, WASHINGTON, DC, vol. 119, no. 22, 4 June 1997 (1997-06-04), pages 5081-5090, XP000775689 ISSN: 0002-7863 * page 5082, scheme 1, compounds 7a - 7k * | 8,9 | |
| X | PIRRUNG ET AL.: "COMPARISON OF METHODS FOR PHOTOCHEMICAL PHOSPHORAMIDITE-BASED DNA SYNTHESIS" JOURNAL OF ORGANIC CHEMISTRY, vol. 60, 1995, pages 6270-6276, XP002088249 * COMPOUNDS 1-4,6, PAGE 6274, COLUMN 1 * | 8,9 | TECHNICAL FIELDS SEARCHED (Int.Cl.7) |
| X | DOUGAN ET AL.: "SYNTHESIS AND RADIOIODINATION OF A STANNYL OLIGODEOXYRIBONUCLEOTIDE" NUCLEIC ACID RESEARCH, vol. 25, no. 14, 1997, pages 2897-2901, XP002167902 * compound 5, page 2899, scheme 1; page 2898, 2nd paragraph, lines 9-11 * | 8,9 | |
| D,X | IWAI ET AL.: "5'-LEVULINYL AND 2'-TETRAHYDROFURANYL PROTECTION FOR THE SYNTHESIS OF OLIGORIBONUCLEOTIDES BY THE PHOSPHORAMIDITE APPROACH" NUCLEIC ACID RESEARCH, vol. 16, no. 20, 1988, pages 9443-9456, XP002167839 * page 9446, table 1 * | 10 | |
| | --- | | |
| | -/-- | | |

EPO FORM 1503 03.82 (P04C10)



European Patent
Office

PARTIAL EUROPEAN SEARCH REPORT

Application Number
EP 99 30 6168

| DOCUMENTS CONSIDERED TO BE RELEVANT | | | CLASSIFICATION OF THE APPLICATION (Int.Cl.7) |
|-------------------------------------|--|-------------------|--|
| Category | Citation of document with indication, where appropriate, of relevant passages | Relevant to claim | |
| P,X | WO 98 39348 A (AFFYMETRIX, INC.) 11 September 1998 (1998-09-11) * page 22, table 1, entries 1,2,5-10; page 19, lines 16-20 * | 1-9 | |
| E | WO 99 54509 A (AFFYMETRIX, INC.) 28 October 1999 (1999-10-28) * 5'-OMeNPOC-(N2-p-isopropylphenoxyacetyl)-2'-deoxyguanosine-3'-O-(2-cyanoethyl)-N,N-diisopropylphosphoramidite, page 40, lines 18-20 * | 8,9 | |
| E | WO 00 61594 A (DEUTSCHES KREBSFORSCHUNGSZENTRUM) 19 October 2000 (2000-10-19) * examples 10-13, 16, page 17 - page 19 * | 8,9 | TECHNICAL FIELDS SEARCHED (Int.Cl.7) |
| E | WO 00 18778 A (PHYLOS, INC.) 6 April 2000 (2000-04-06) * example 2a, page 16 * | 8,9 | |
| E | US 6 022 963 A (MCGALL ET AL.) 8 February 2000 (2000-02-08) * column 13, example * | 8,9 | |

EPO FORM 1503 01.82 (P04C10)

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 99 30 6168

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

28-05-2001

| Patent document cited in search report | Publication date | Patent family member(s) | Publication date |
|---|---------------------|----------------------------|---------------------|
| WO 9628457 A | 19-09-1996 | US 5908926 A | 01-06-1999 |
| | | AU 5422896 A | 02-10-1996 |
| | | CA 2215657 A | 19-09-1996 |
| | | EP 0815114 A | 07-01-1998 |
| | | JP 11501936 T | 16-02-1999 |
| EP 0742287 A | 13-11-1996 | US 6156501 A | 05-12-2000 |
| WO 9839348 A | 11-09-1998 | US 6147205 A | 14-11-2000 |
| WO 9954509 A | 28-10-1999 | AU 3659199 A | 08-11-1999 |
| | | EP 1071821 A | 31-01-2001 |
| WO 0061594 A | 19-10-2000 | DE 19915867 A | 19-10-2000 |
| | | AU 5059800 A | 14-11-2000 |
| WO 0018778 A | 06-04-2000 | AU 6270499 A | 17-04-2000 |
| US 6022963 A | 08-02-2000 | US 6147205 A | 14-11-2000 |

EPO FORM P453

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

THIS PAGE BLANK (USPTO)